

## ABSTRACT

### Background

Induction of labor is being increasingly used to prevent adverse outcomes in the mother and the newborn. This study assessed the prevalence of induction of labor and determinants of its use in Africa.

### Methods

We performed secondary analysis of the WHO Global Survey of Maternal and Newborn Health of 2004 and 2005. The African database was analyzed to determine the use of induction of labor at the country level and indications for induction of labor. The un-met needs for specific obstetric indications and at country level were assessed. Determinants of use of induction of labor were explored with multivariate regression analysis.

### Results

A total of 83,437 deliveries were recorded in the 7 participating countries. Average rate of induction was 4.4% with a range of 1.4 to 6.8%. Pre-labor rupture of membranes was the commonest indication for induction of labor. Two groups of women were identified: 2,776 women with indications had induction of labor while 7,996 women although had indications but labor was not induced.

Induction of labor was associated with reduction of stillbirths and perinatal deaths [OR 0.34; 95% CI (0.27 to 0.43)].

Unmet need for induction of labor ranged between 66.0% and 80.2% across countries.

Determinants of having an induction of labor were place of residence, duration of schooling, type of health facility and level of antenatal care.

### Conclusion



*Your complimentary  
use period has ended.  
Thank you for using  
PDF Complete.*

[\*Click Here to upgrade to  
Unlimited Pages and Expanded Features\*](#)

h facilities in Africa is very low. Improvements in social  
reverse the high unmet need for induction of labor.