

**ANTIMICROBIAL ACTIVITY, TOXICITY AND PHYTOCHEMICAL ANALYSIS
OF FOUR MEDICINAL PLANTS TRADITIONALLY USED IN MSAMBWENI
DISTRICT, KENYA**

MAGRATE MUTHONI KAIGONGI,

B.Sc. (Biology)

(Nairobi).

**A THESIS SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS
FOR THE DEGREE OF MASTER OF SCIENCE IN PLANT TAXONOMY AND
ECONOMIC BOTANY OF THE UNIVERSITY OF NAIROBI**

**SCHOOL OF BIOLOGICAL SCIENCES,
UNIVERSITY OF NAIROBI.**

August, 2014.

DECLARATION

This thesis is my original work and has not been presented for a degree in any other University.

Magrate Muthoni Kaigongi, BSc.
School of Biological Sciences,
College of Biological and Physical Sciences,
University of Nairobi

Signed : _____ Date: _____

Supervisors

This thesis has been submitted for examination with our approval as University supervisors

Dr. Dossaji Saifuddin F., BSc, MSc, PhD.
School of Biological Sciences,
College of Biological and Physical Sciences,
University of Nairobi

Signature: _____ Date: _____

Dr. Joseph Mwanzia Nguta, BVM, MSc, Ph.D. (UON).
Department of Public Health, Pharmacology and Toxicology,
Faculty of Veterinary Medicine,
College of Agriculture and Veterinary Sciences,
University of Nairobi

Signature: _____ Date: _____

Dr. Catherine Lukhoba W. BSc, MSc, PhD.
School of Biological Sciences,
College of Biological and Physical Sciences,
University of Nairobi

Signature: _____ Date: _____

DEDICATION

This thesis is dedicated to my family members.

ACKNOWLEDGEMENTS

First, I am very grateful to the Almighty God for the wisdom, guidance and good health that He has accorded to me throughout my studies. I also express my appreciation to my

supervisors; Drs. Saifuddin Dossaji, Joseph Nguta and Catherine Lukhoba for their suggestions, guidance, encouragement and support throughout the period of this study.

I would also like to take this opportunity to thank Ms. Trizah Wangechi for taking me through microbiological techniques and making herself available at all times to solve all my problems on microbiological work. Thank you, Mr. Ndi and Mr. Wachira for your help in the microbiology lab.

It is impossible to frame the help of Ms. Judith Odhiambo in words; whether I needed laboratory materials or some form of advice, you were always there for me. The greatest way to say thank you to Judith would be to learn some of your very good habits. I surely want to emulate you.

To Mr. Fredrick Musila, thank you for taking me through the brine shrimp lethality technique. To my fellow graduate students, Ms. Jacklyne Cherotich and Mr. Victor Sirama, thank you for your help and support throughout this study.

I want to thank the Masinde Muliro University of Science and Technology for awarding me the scholarship which has provided me an opportunity to pursue my studies. To University of Nairobi, thank you for offering me space and facilities to carry out this study.

I wish to extend my final thanks to my family, especially my sister Lucy and my brother Muriungi, without whose love and support, I would not be here. Thank you very much for being by my side, believing in me, and putting up with me during the good and bad times. Once again, thank you.

TABLE OF CONTENTS

DECLARATION	i
DEDICATION.....	iii
ACKNOWLEDGEMENTS.....	iii
TABLE OF CONTENTS	v
LIST OF TABLES	vii
LIST OF FIGURES	viii
LIST OF APPENDICES	ix
LIST OF ABBREVIATIONS.....	x
ABSTRACT	xi
CHAPTER ONE: INTRODUCTION.....	1
CHAPTER TWO: LITERATURE REVIEW.....	3
2.1. Microbial infections and their health effects.....	3
2.2. Methicillin (ATCCMP-2) resistant <i>Staphylococcus aureus</i> (MRSA)	3
2.3. <i>Bacillus cereus</i>	4
2.4. <i>Escherichia coli</i>	4
2.5. <i>Pseudomonas aeruginosa</i>	5
2.6. <i>Candida albicans</i>	5
2.7. Antibiotic resistance and its socio-economic impacts	6
2.8. Traditional medicine in Africa	6
2.9. Conventional antibiotics and the problem of microbial resistance.....	7
2.10. Plant derived antimicrobials	8
2.11. Plants systematics, phytoconstituents and their economic importance	10
2.11.1. <i>Adansonia digitata</i> L. (Bombacaceae)	10
2.11.2. <i>Launaea cornuta</i> (Hocht. ex Oliv. & Hern) C.Jeffrey (Compositae).....	12
2.11.3. <i>Grewia trichocarpa</i> Hochst. ex A.Rich (Tiliaceae)	14
2.11.4. <i>Zanthoxylum chalybeum</i> Engl. (Rutaceae)	15
2.12. Toxicity Studies	18
2.13. Significance of toxicity studies.....	19
2.14. Brine shrimp lethality assay	20
2.15. Clinical and Laboratory Standards Institute protocol on evaluation of antimicrobial activity	21

2.16. Interpretation of antimicrobial results.....	21
2.17. Problem statement.....	22
2.18. Justification	23
2.19. Objectives.....	23
2.19.1. Main Objective.....	23
2.19.2. Specific objectives.....	23
2.19.3. Null Hypothesis	24
CHAPTER THREE: MATERIALS AND METHODS.....	25
3.1. Collection of plant material.....	25
3.2. Preparation of Crude Extracts	25
3.3. Microorganism Suspension Standardization.....	26
3.4. Disc Diffusion Test.....	26
3.5. Determination of minimum inhibitory concentration (MIC)	27
3.6. Acute toxicity testing.....	28
3.7. Thin Layer Chromatography (TLC)	29
3.8. Data analysis.....	30
CHAPTER FOUR: RESULTS	32
4.1 Extraction of collected plants.....	32
4.2. Antimicrobial activity	33
4.2.1. Disc Diffusion Technique.....	33
4.2.2. Broth dilution technique	39
4.3. Brine-Shrimp lethality test	41
4.4. Phytochemical screening of secondary compounds	42
CHAPTER FIVE: DISCUSSION.....	44
CHAPTER SIX: CONCLUSIONS AND RECOMMENDATIONS.....	52
REFERENCES	53
APPENDICES	63

LIST OF TABLES

Table 1: Selected plant species from Msambweni district used in traditional medicine	18
Table 2: Plants Collected from Msambweni District, Kenya.....	25
Table 3: List of microbes tested in the study.....	26
Table 4: Procedure followed in screening for different chemical compounds.....	30
Table 5: Percentage yield of plant extracts using aqueous and organic extraction methods ..	32
Table 6: Minimum Inhibitory Concentration (MIC) of Organic Crude Extracts	40
Table 7: Minimum Inhibitory Concentration (MIC) of Aqueous Crude Extracts.....	41
Table 8: LC ₅₀ values for various tested aqueous crude extract on Brine-shrimp larvae	41
Table 9: LC ₅₀ values for various tested organic crude extract on Brine-shrimp larvae.....	42
Table 10: Presence of alkaloids, flavonoids, sesquiterpene- lactones and saponins in aqueous extracts.....	43
Table 11: Presence of alkaloids, flavonoids, sesquiterpene- lactones and saponins in organic extracts.....	44

LIST OF FIGURES

Figure 1: <i>Adansonia digitata</i> tree (Bombacaceae).....	12
Figure 2: A flowering herb of <i>Launaea cornuta</i> (Compositae).....	14
Figure 3: <i>Grewia trichocarpa</i> (Tiliaceae) showing leafy shoot	15
Figure 4: <i>Zanthoxylum chalybeum</i> (Rutaceae) tree	17
Figure 5: Inhibition zones caused by <i>Z. chalybeum</i> at 200 mg/ml and positive control against <i>B. cereus</i>	34
Figure 6: Growth inhibition of <i>B. cereus</i> by different concentrations of extracts from the four test plants	35
Figure 7: Growth inhibition of <i>P. aeruginosa</i> by different concentrations of extracts from the four test plants.....	36
Figure 8: Growth inhibition of MRSA by different concentrations of extracts from the four test plants	37
Figure 9: Inhibition zones showing antifungal activity of <i>A. digitata</i> at 200 mg/ml and positive control against <i>C. albicans</i>	38
Figure 10: Growth inhibition of <i>C. albicans</i> by different concentrations of extracts from the four test plants.....	39
Figure 11: A photograph showing sesquiterpene lactones results of the tested extracts on a TLC plate.....	42

LIST OF APPENDICES

Appendix 1: Growth inhibition of <i>B. cereus</i> by aqueous extracts.....	63
Appendix 2: Growth inhibition of <i>B. cereus</i> by organic extracts.....	64
Appendix 3: Growth inhibition MRSA by aqueous extracts.....	65
Appendix 4: Growth inhibition of MRSA by the organic extracts.....	66
Appendix 5: Growth inhibition of <i>P. aeruginosa</i> by aqueous extracts.....	67
Appendix 6: Growth inhibition of <i>P. aeruginosa</i> by organic extracts.....	68
Appendix 7: Growth inhibition of <i>C. albicans</i> by organic extracts.....	69
Appendix 8: Brine shrimp lethality raw data.....	70

LIST OF ABBREVIATIONS

ANOVA:	Analysis of Variance
ATCC:	American Type Culture Collection
BSLB:	Brine Shrimp Lethality Bioassay
CMR:	Centre for Microbiology Research
CLSI:	Clinical and Laboratory Standards Institute
CHCl ₃ :	Chloroform
DMSO	Dimethylsulphoxide
DPHPT:	Department of Public Health, Pharmacology and Toxicology
H ₂ SO ₄ :	Sulphuric acid
KEMRI:	Kenya Medical Research Institute
LC ₅₀ :	Lethal Concentration 50
MIC:	Minimum Inhibitory Concentration
MRSA:	Methicillin Resistant <i>Staphylococcus aureus</i>
MeOH:	Methanol
SPSS:	Statistical Package for the Social Sciences
TLC:	Thin Layer Chromatography
UoN:	University of Nairobi
UV :	Ultra Violet Light
WHO:	World Health Organization

ABSTRACT

Introduction: Antimicrobial resistance to currently utilized drugs has caused major challenges in both the agricultural and medical sectors in the past years. This resistance has not only hampered the treatment of microbial diseases, but has also increased the cost of treatment which is especially burdening third world countries. Therefore, there is need to find alternative methods of curbing the infectious microorganisms.

Methodology: The current study was designed to evaluate the antimicrobial activity, toxicity and phytochemical composition of organic and aqueous crude extracts from each plant were evaluated for their *in vitro* antimicrobial activity against methicillin resistant *Staphylococcus aureus* (MRSA), *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans* using broth dilution and disc diffusion methods. Toxicity was determined using Brine-shrimp larvae assay while the presence of flavonoids, alkaloids, saponins and sesquiterpene lactones was determined using Thin Layer Chromatography (TLC). Analysis of variance (ANOVA) was used to analyse the disc diffusion results. Finney's computer programme was used to determine the LDC₅₀ of each extract.

Results: It was observed that the organic crude extracts of all the tested plants except *L. cornuta* showed activity against four micro-organisms and at concentrations lower than the aqueous crude extracts. None of the crude extracts elicited any inhibition against *E. coli*. *Adansonia digitata* and *G. trichocarpa* had LC₅₀>1000 µg/ml and were shown to be non-toxic to Brine shrimp larvae unlike those of *Z. chalybeum* and *L. cornuta* which both had LC₅₀<500 µg/ml and were considered to be toxic. Phytochemical screening of the crude extracts showed that alkaloids, flavonoids, sesquiterpene lactones and saponins were present in the four plants under investigation.

Conclusion: This study provides the first record of antimicrobial activity, toxicity and phytochemical composition of *G. trichocarpa*. The study has shown that *A. digitata* and *Z. chalybeum* possess promising antimicrobial activity against microbes of health importance and could lead to the isolation of new and effective antimicrobial compounds. Further research should be carried on *Z. chalybeum* and *A. digitata* to isolate and characterize the compounds responsible for the observed activity.

Key words: Antimicrobial activity, *Adansonia digitata*, *Launaea cornuta*, *Zanthoxylum chalybeum*, *Grewia trichocarpa*, Brine shrimp lethality test, Phytochemical analysis,

CHAPTER ONE: INTRODUCTION

A two-fold increase in microbial resistance against antibiotics has developed over the last forty years in both medical and livestock sector (Daboor and Haroon, 2012). Microbial infections have caused a big burden of diseases and bacteria are listed in the first position among common microorganisms responsible for opportunistic diseases associated with HIV/AIDS (Rathee *et al.*, 2012). Increased antibiotic resistance has become a global concern, coupled with the problem of microbial persistence, thus highlighting the need to develop novel microbial drugs that are not only active against drug resistant microbes, but more importantly, kill persistent micro-organisms and shorten the length of treatment (Mariita *et al.*, 2010). People in various parts of Africa have since time immemorial have used plant extracts to treat themselves together with their livestock whenever they became sick or injured in their environment (Kokwaro, 2009). According to World Health Organisation (2001), 80% of African and Asian populations depend on traditional medicine particularly plant biodiversity for primary health care because plant derived medicines are relatively cheaper and safer compared to the synthetic alternatives. Many commercial drugs used in modern medicine are derived from plants following ethno-botanical and ethno-medical knowledge (Arokiyaraj *et al.*, 2012). Medicinal plants have bioactive compounds which treat various ailments caused by microorganisms. These compounds may have evolved in plants as self defence against pests and pathogens to help plants to establish themselves in their environment (Sukumaran *et al.*, 2011).

The World Health Organisation (WHO, 2001) has advocated the use of traditional medicine as a safe remedy for ailments of microbial and non-microbial origin. Plant derived secondary metabolites such as alkaloids, flavonoids, phenolics, saponins, and tannins protect plants from invaders such as fungi, bacteria, viruses, and nematodes. It is estimated that among the traditional medicinal plants, only 12,000 or less than 10% have undergone pharmacological

evaluation (Marimuthu *et al.*, 2011). It is therefore possible that many plants not yet validated may be the source of drugs that may be effective against resistant pathogens to avoid the threat of post biotic era (Marimuthu *et al.*, 2011). The importance of a country's diverse medicinal plants lies not only in its chemotherapeutic value in traditional medicine but also in its potential in drug discovery and this prompts a need to research on Kenyan medicinal plants (Vital and Rivera, 2011). Traditional medicine is widely practiced in Kenya and about 400 plant species have been recorded to be used in traditional remedies (Kokwaro, 2009). In the rural areas, reliance on traditional medicine is high and is attributed to both economic and cultural factors (Aketch, 1992).

Plants have toxic effects on livestock and humans, such toxicity may lead to death of an animal or humans consuming the plant (Hood, 2009). In traditional medicine, there is likelihood to overdose the patient due to imprecise nature of diagnosis and dosage not only in Kenya but worldwide. This is not only unique to traditional medicine but can also occur in modern medicine (Kokwaro, 2009). Toxicity is attributed to certain active principles found in plants; these chemical substances interact with living systems and affect normal processes. All chemicals can cause harm to organisms at some level of exposure. Toxicity tests are important so as to determine the lethality of drugs and to determine the harmless concentration of drugs for consumption (Hood, 2009). Biological activity of plants is attributed to the class and concentration of phytochemical constituents which makes some plant extracts exhibit a variety of activities (Wang *et al.*, 2010). Phytochemical screening of plant extracts especially those which have been used in traditional medicine is therefore essential so as to identify phytochemical constituents in the plants that are responsible for a given bioactivity. The current study investigated the antimicrobial activity, toxicity and phytochemical composition of *Adansonia digitata*, *Zanthoxylum chalybeum*, *Launaea cornuta* and *Grewia trichocarpa* traditionally used in Msambweni ethnomedicine.

CHAPTER TWO: LITERATURE REVIEW

2.1. Microbial infections and their health effects

Microbial infections have caused a big burden of diseases and bacteria are listed in the first position among common microorganisms responsible for opportunistic diseases associated with HIV/AIDS (Rathee *et al.*, 2012). In developing countries, bacterial infections are prevalent due to factors such as poor hygiene, sanitation and overcrowding in the living conditions (Rosoanaivo and Ratsimamanga-Urveg, 1993). Throughout history, infectious diseases have been a major threat to human and animal health and a prominent cause of morbidity and mortality, WHO, (2003). Increased antibiotic resistance has become a global concern, coupled with the problem of microbial persistence, thus highlighting the need to develop novel microbial drugs that are not only active against drug resistant microbes, but more importantly, kill persistent micro-organisms and shorten the length of treatment. Apart from toxicity, lengthy therapy also creates poor patient compliance (Mariita *et al.*, 2010).

2.2. Characteristics of Various Microbial Pathogens

2.2.1. Methicillin (ATCCMP-2) resistant *Staphylococcus aureus* (MRSA)

Staphylococcus aureus is a facultative anaerobic Gram positive bacterium belonging to the family Staphylococcaceae and is frequently found in the human respiratory tract and on the skin. It is a non motile and non spore forming bacteria that appears in grape-like clusters with large, yellow colonies and reproduces asexually by binary fission (Nizet and Bradley, 2011). It is a versatile organism and a source of a wide range of infectious diseases in humans such as life-threatening deep seated infections like bacteremia, pneumonia and meningitis (Kanafani and Fowler Jr., 2006). In burn patients, wound sepsis remains a major cause of death due to colonisation with methicillin resistant *S. aureus* on the surface of burn wounds which may go on to cause systemic infections and other serious clinical complications. *S. aureus* resistance

against methicillin was first reported in 1960s soon after introduction of penicillinase-resistant β -lactam antibiotics methicillin (Cook *et al.*, 1998). In addition to their resistance against all β -lactam antibiotic, methicillin resistant *S.aureus* strains are also resistant to several other classes of antibiotics including the Aminoglycosides, Quinolones, Clindamycin and Erythromycin. There are several mechanisms responsible for the resistance, the major one being production of penicillin-binding protein *2a* encoded by *mecA* gene, two additional mechanisms are hyper-production of penicillinase and modified affinity of usual penicillin-binding proteins (Durmaz *et al.*, 1997).

2.2.2. *Bacillus cereus*

B. cereus is a Gram positive, endospore-forming, facultative, rod shaped motile bacterium (Pirtijärvi *et al.*, 1999). It is found widely distributed in nature including mucous membrane and faeces of healthy humans (Williams, 1986). *B. cereus* causes food poisoning of two types characterized by abdominal pain and diarrhea for 8-16 hours or by nausea and vomiting 1-5 hrs after ingestion of contaminated food (Asano *et al.*, 1997; Gilbert, 1979). Infection of central nervous system is rare but it is associated with immunosuppression and central nervous system invasive devices (Drobniewski, 1993). Some *B. cereus* isolates according to Waber *et al.* (1988) are susceptible *in vitro* to vancomycin, gentamicin, clindamycin, erythromycin and chloramphenicol.

2.2.3. *Escherichia coli*

It is a facultative anaerobic, Gram- negative, non- spore forming, rod-shaped bacterium that can be a commensal or a deadly pathogen in the lower intestine of warm blooded organisms (Kaper *et al.*, 2004). Enteropathogenic *E. coli* causes watery diarrhea which in serious cases resemble classic cholera among infants (Cravioto *et al.*, 1991). According to Čížek *et al.*

(2008), some isolates are resistant to chloramphenicol, sulphonamide and trimethoprim-sulphomethoxale combination.

2.2.4. *Pseudomonas aeruginosa*

It is a Gram- negative bacterium occurring as short chains, pairs or singly with an optimum growth temperature of 37⁰c. It is widely found in moist environment with low nutrient and ion concentration (Bennik *et al.*, 1999). *Pseudomonas aeruginosa* is an opportunistic pathogen causing serious infections to already damaged tissues such as burns or to immunosuppressed individuals such as cancer and HIV/AIDS patients (McCarthy, 1990; Soltan,1998). *P. aeruginosa* is one of the most resistant microbes encountered in clinical laboratories (Shimeld and Rodgers, 1999).

Some strains such as metallo-beta-lactamase producing are reported to be resistant to all antibiotics except colistin (Edelstein *et al.*, 2013).

2.2.5. *Candida albicans*

Candida albicans is a diploid, dimorphic, unicellular yeast causing opportunistic oral (Thrush disease) and genital (Candidiasis) infections in humans (Pesti *et al.*, 2012). Oral infection caused by *C. albicans* is higher in diabetic mellitus patients due to high sugar concentrations (Mubarak *et al.*, 2013). It is believed to be one of the fungi that play a role in sexually transmitted diseases especially in immune-compromised and HIV/AIDS patients (*et al.*, 2011).

Studies have shown that oral candidiasis, mostly commonly characterized by development of oral thrush, is the most frequent AIDS-associated opportunistic infection, with up to 90% of HIV-infected individuals suffering at least one episode during the course of their disease

(Vazquez, 1999). *C. albicans* is reported to be resistant against Fluconazole, Flucytosine, and intraconazole whereas resistance to AmphotericinB is rare (Marchese *et al.*, 2007).

2.3. Antibiotic resistance and its socio-economic impacts

Millions of lives have been saved due to substantial reduction in mortality achieved with antibiotic therapy. However, the antibiotic effectiveness almost all the available antibiotics is being threatened by the rising resistance of microorganisms. The major cause of this resistance is the over-use and misuse of antibiotics which is exerting undue selective pressure on microorganisms (Paphitou, 2013). This has made infections more difficult to treat which is contributing to the high morbidity and mortality of previously treatable infections (Byarugaba, 2004).

Infections caused by antibiotic resistant organisms are more likely to prolong hospitalisation, increase the risk of death and require more toxic and expensive antibiotics (Graf and Martin, 2000). In addition, as the prevalence of multidrug resistant organisms increase, these additional costs will become a greater threat to the local, regional and national medical care systems, many of which are already struggling to survive (DiazGranados *et al.*, 2008). The pharmaceutical industry finds it increasingly difficult to keep pace with the antimicrobial resistance and is no longer reliable in bringing novel and more effective drugs to the market (Graf and Martin 2000).

2.4. Traditional medicine in Africa

The use of natural products with therapeutic properties is as ancient as human civilization and for a long time, mineral, plant and animal products were the main source of drugs in Africa (Maryam *et al.*, 2010). About 80% of the world's people rely on traditional plant based medicine (UNEP, 2010). It is estimated that about 75% population in Kenya seeks health care

among traditional healers (Sandiga *et al.*, 1995). Traditional medicine is widely practiced in Kenya, and has been documented by various ethnomedical surveys (Miaron *et al.*, 2004; Kareru *et al.*, 2007).

The high cost of important conventional drugs and/ or inaccessibility to western health care facilities has led to over-reliance on traditional medicine since it is affordable and available to people. On the other hand, even when western health facilities are available, traditional medicine is viewed as an efficient and an acceptable system from a cultural perspective (Munguti, 1997). Infections associated with bacterial and fungal pathogens are among some of the indications treated using traditional remedies in Kenya (Njoroge and Bussmann, 2007).

2.5. Conventional antibiotics and the problem of microbial resistance

Conventional antibiotics refer to the synthetic chemicals used as bactericides and fungicides. Antibiotics act in various ways such as; inhibition of cell membrane functions, inhibition of protein synthesis, inhibition of nucleic acid synthesis and inhibition of cell wall synthesis (Kalayou *et al.*, 2012). Since their discovery, antimicrobial drugs have proved remarkably effective for the control of bacterial infections. However, it was soon evident that bacterial pathogens were unlikely to surrender unconditionally, because some pathogens rapidly become resistant to many of the first discovered effective drugs (Cowan, 1999).

Due to indiscriminate use of antimicrobial drugs, microorganisms have developed resistance to many antibiotics and that has created immense clinical problems in the treatment of infectious diseases (Davis, 1994). In addition, antibiotics are associated with adverse effects on host, which include depletion of beneficial gut and mucosal microorganisms, immunosuppression, hypersensitivity and allergic reactions. The drug resistant bacteria have further complicated the treatment of infectious diseases in immunocompromised, AIDS and cancer patients especially in the case of nosocomial infections (McGaw *et al.*, 2001). There is not only the lost of

effectiveness of antibiotics against multidrug resistant microorganisms, but also global problem for the lost of budget for treating infectious diseases (Ahmad and Beg, 2001).

The emergence of antimicrobial resistance has its roots in the use of antimicrobials in animals and the subsequent transfer of resistance genes and bacteria among animals, animal products and the environment (McEwen and Fedorka-Cray, 2002). Extra-chromosomal genes were found responsible for these antimicrobial resistant phenotypes that may impart resistance to an entire antimicrobial class. These resistance genes have been associated with plasmids which are large, transferable, extra-chromosomal DNA elements. Other DNA mobile elements, such as transposons and integrons, are present on plasmids. These DNA mobile elements transmit genetic determinants for antimicrobial resistance mechanisms and may cause rapid dissemination of resistance genes among different bacteria (Toutou *et al.*, 2004). The emergence of multiresistant bacteria to antimicrobial drugs has increased the need for new antibiotics or modifications of older antibiotics (Tollefson and Miller, 2000).

2.6. Plant derived antimicrobials

The antimicrobial compounds from plants may inhibit bacterial growth by different mechanisms than those presently used antimicrobials and may have a significant clinical value in treatment of resistant microbial strains (Shankar *et al.*, 2010). Extracts isolated from several plants have been reported to have biological activity such as antimicrobial, anti-inflammatory and antioxidant activities (Yusuf *et al.*, 2001).

Phenolic compounds are one of the most diverse groups of secondary metabolites found in edible plants. They are found in a wide variety of fruits, vegetables, nuts, seeds, stems and flowers as well as tea, wine, propolis and honey, and represent a common constituent of the human diet. In nature they are involved in plant growth and reproduction, provide resistance from pathogens and predators and protect crops from disease and pre-harvest seed germination

(Ross and Kasum 2002). There are different classes of polyphenols known as tannins, lignins and flavonoids. Each class of polyphenols possesses chemical characteristics that set them apart from one another.

Flavonoids are the most widely occurring polyphenol and are present in almost every form of human consumed vegetation. Dietary flavonoids have attracted interest because they have a variety of beneficial biological properties, which may play an important role in the maintenance of human health. Flavonoids are potent antioxidants, free radical scavengers and metal chelators; they inhibit lipid peroxidation and exhibit various physiological activities including anti-inflammatory, anti-allergic, anti-carcinogenic, anti-hypertensive, anti-arthritic and antimicrobial activities. Consumption of phenol-rich beverages, fruit and vegetables has commonly been associated with reduction of the risk of cardiovascular diseases in epidemiological studies (Whiting 2001). Flavonoids have been found to be the most abundant polyphenols. The biosynthesis of flavonoids is stimulated by sunlight (ultraviolet radiation), so higher concentrations of flavonoids can typically be found in the outer most layers of fruits and vegetables (i.e. the skins). Flavonoids can be divided into six subclasses according to the degree of oxidation of the oxygen heterocycle: flavones, flavonols, isoflavones, anthocyanins, flavanones and flavonols (catechins and anthocyanidins).

Extraction of polyphenols can be performed using a solvent like water, hot water, methanol, methanol/formic acid, methanol/water/acetic or formic acid etc. Therefore, the total polyphenol amounts detected from the same plant and their corresponding antioxidant and antimicrobial activities may vary widely, depending on external conditions applied. It was reported that an antimicrobial action of phenolic compounds was related to inactivation of cellular enzymes, which depended on the rate of penetration of the substance into the cell or caused by membrane permeability changes (Schultz *et al.*, 1992).

Terpenoids and essential oils are also plant derived antimicrobials. The oils are secondary metabolites which are highly enriched in compounds based on isopropene structure. They are called terpenes. When compounds contain additional elements usually oxygen, they are termed terpenoids (Singh and Handique, 1997). Terpenes or terpenoids are active against bacteria, viruses and protozoa. The mechanism of action of terpenes is not fully known but it is speculated to involve membrane disruption by lipophilic compounds (Hamed, 2011).

Heterocyclic nitrogen compounds are called alkaloids. The mechanism of action of highly aromatic planar quaternary alkaloids is attributed to their ability to intercalate with DNA of microorganisms (Cowan, 1999).

Lectins and polypeptides are often positively charged and contain disulphide bonds. Their mechanism of action may be the formation of ion channels in the microbial membrane or competitive inhibition of adhesion of microbial proteins to host polysaccharide receptors (Sher, 2004)

2.7. Plants systematics, phytoconstituents and their economic importance

There are several approaches for selecting plant species for biological investigation. Selection of species based on their use in traditional medicine is reported to be more valuable and many drugs that are of plant origin were discovered from plants used in traditional medicine (Cotton, 1996).

2.7.1. *Adansonia digitata* L. (Bombacaceae)

Adansonia digitata, is commonly known as baobab, is a tree that grows in dry parts of Africa. It has leaves in form of five-leaflets, large showy white flowers and the fruit is in form of a pulp. It is a multipurpose tree with leaves eaten as fresh vegetable, dried pulp used to prepare juice and different parts utilised in cure of various diseases. The pulp, leaves and seeds contain

calcium, magnesium, potassium, zinc, iron and vitamins A and C in variable proportions (Assogbadjo *et al.*, 2008).

Proanthocyanidins, terpenoids, flavonoids, and sterols have been isolated from *A. digitata* (VanStaden *et al.*, 2012). Compaoré *et al.* (2011) has shown that flavonoids, proanthocyanidins and phenolic compounds are present in the pulp of *A. digitata* which make it a good radical scavenger. Elsewhere the fruits of *A. digitata* have been reported to contain proanthocyanidins as the major compounds (Shahat, 2006). Triterpenes, alkaloids, anthraquinones, saponins, tannins have also been reported to be present in the fruit pulp of *A. digitata* (Ramadan *et al.*, 1994; Gbadamosi *et al.*, 2011).

The plant has anti-inflammatory, antipyretic, antimicrobial and analgesic activities (Ramadan *et al.*, 1994) Powdered leaves have anti-asthmatic and have anti-tension and antihistamine properties. Leaves are also used for other conditions such as diarrhea, dysentery, ophthalmia and otitis media. Bark is used as a substitute for quinine to curb high fever and act as prophylactic measure for malaria as well (Sidibe and Williams, 2002). The stem extracts are non-toxic to brine shrimp larvae and aqueous extract has a high anti-malarial activity (Musila *et al.*, 2013). In other studies VanStaden *et al.* (2011), *A. digitata* has showed considerable antimicrobial activity against *Bacillus subtilis*, *S. aureus* and *C. albicans*.



Figure 1: *Adansonia digitata* tree (Bombacaceae)

(Photograph: Magrate Kaigongi in Tharaka-Nithi County)

2.7.2. *Launaea cornuta* (Hocht. ex Oliv. & Hern) C.Jeffrey (Compositae)

L. cornuta is a perennial herb with erect stem, rosette of leaves and a diffuse inflorescence growing in disturbed and artificial grasslands and roadsides. The fresh young leaves and shoots are used as vegetable as the plant contains vitamin C, sodium, potassium, calcium and iron (Katariina, 2000). Lyimo *et al.* (2003) also reported that the leafy vegetable contains vitamin C, iron, fat, calcium and protein. A decoction from the whole plant is used to treat cancer and as a remedy to diabetes (Kareru *et al.*, 2007). Leaves and roots of *L. cornuta* are used as antimalarial, antimicrobial and used by the Suba people in Kenya to treat opportunistic diseases associated with HIV/AIDS. They also treat stomach ache and stop blood flow during circumcision (Nagata *et al.*, 2011).

The genus *Launaea* is characterized by flavonoids, triterpenes, sesquiterpene lactones, coumarins and steroids (Ali *et al.*, 2003). Flavones like apigenin and luteolin are common compounds in the genus. Steroids like stigmasterol, cholesterol, taraxasterol have been identified in *L. nudicaulis*. Flavone glycosides such as apigenin-7-glycoside, luteolin-7-glycoside, luteolin-7-rutinoside and Vitexin have been isolated from the extract of *L. tenuiloba* and *L. resedifolia* an anthocyanin is found in *L. asplenifolia* (Ali *et al.*, 2003; Fairouz *et al.*, 2010).

Related species that have antimicrobial activity include: *L. cassiniana* Kuntze has nematocidal activity, *L. intybacea* likewise was found to have hepatoprotective activity on paracetamol introduced hepatotoxicity in albino rats (Takate *et al.*, 2010) and *L. sonchoides* inhibits the growth of the fungi *Trichoderma hamatum* and *T. viridae* (Abou-Zeid *et al.*, 2008).

Various species of *Launaea* such as *L. arborescens*, *L. mucronata*, *L. nudicaulis* and *L. capitata* contains several types of flavones such as luteolin, apigenin, and flavone glycosides such as apigenin 7-O-glucoside, vitexin, luteolin 7-O-glucoside and luteolin 7-O-rhamnoside (Christian and Vipaporn, 2010). Two isoprenylated flavonoids; asplenitin and asplenetin 5-O-neohesperidoside have been identified in *L. asplenifolia* (Denis and Ragaj, 1996). Roots of *L. mucronata* Muschl, aerial parts of *L. spinosa* Sch and aerial parts and roots of *L. tenuiloba* contain lactucin type guianolides as stated by Christian (2008).

The crude extracts (both organic and aqueous extracts) have moderate antimalarial activity and highly toxic to the Brine-shrimp larvae. They also contain flavonoids, alkaloids and saponins with sesquiterpene lactone being only present in the organic extract (Musila *et al.*, 2013)



Figure 2: A flowering herb of *Launaea cornuta* (Compositae)

(Photograph: Magrate Kaigongi in Tharaka-Nithi County)

2.7.3. *Grewia trichocarpa* Hochst. ex A.Rich (Tiliaceae)

It is an evergreen shrub or small tree growing up to 6m The Leaves are alternate, simple; stipules subulate, 6–8 mm long. The flowers are yellow to orange with superior ovary. The fruit is drupe (Beentje, 1994). It is used in various ways; the bark fibre is used in hut construction and in basketry. The pounded bark is used for soap-making. The woody branches and wood are used for withies, construction, floors of granaries, tool handles, walking sticks, bows and spear- and arrow shafts. It is also used as fuel wood. The aerial parts are used as fodder. The flowers are a source of bee forage. The ripe fruit is eaten (Beentje, 1994). No

available literature on antimicrobial activity, toxicity and phytochemistry reports on *G. trichocarpa* exists.

Grewia trichocarpa is used among the Luo community to clean snake bite wounds in both humans and livestock (Kokwaro, 2009). *Grewia hexaminta*, one cup is taken 3 times daily for 3-4 days among the Msambweni people of Kenya to treat malaria (Nguta *et al.*, 2011) and has Triterpenoid compounds (Raghunathaiyar, 1996). No available reports on the phytoconstituents of *G. trichocarpa* exist.



Figure 3: *Grewia trichocarpa* (Tiliaceae) showing leafy shoot

(Photograph: Magrate Kaigongi in Kilifi County)

2.7.4. *Zanthoxylum chalybeum* Engl. (Rutaceae)

It is a deciduous tree growing up to 12m high with pale grey bark, dark scales and prickles and a rounded crown. It has compound leaves, usually 3-5 pairs of shiny leaflets with a strong citrus smell when crushed. Flowers are sweet scented, inconspicuous, yellowish-green, produced immediately below the leaves at the base of the new branchlets. The fruits are

spherical (Beentje, 1994). It grows in low altitude areas, dry woodlands and savannah grasslands. Leaves are used as fodder and can be brewed to make tea while the bark is a source of durable timber and charcoal (Bamford and Henderson, 2003).

Leaf decoction is rubbed on swelling for the treatment of oedema in kwashiorkor and snake bites, root decoction is drunk for the treatment of chest pain, pneumonia and bark decoction drunk for the treatment of malaria, colds, coughs, dizziness and also chewed for toothache (Kokwaro, 2009). Seed extracts of *Z. chalybeum* has antiviral activity against measles virus (Olila *et al.*, 2002). The plant has been used traditionally in the management of diabetes (Keter and Mutiso, 2011). *Z. chalybeum* has shown *in vitro* antiplasmodial activity (Rukunga *et al.*, 2009; Nguta *et al.*, 2010), antifungal activity against *Penicillium crustosum* and *Saccharomyces cerevisiae* (Taniguchi *et al.*, 1978), antibacterial activity against *Staphylococcus aureus* (Matu and vanStaden, 2003) and cytotoxic activity against Brine-shrimp larvae (Nguta *et al.*, 2011). *Zanthoxylum rhifolium* is used in treating headaches, anaemia, jaundice and high blood pressure (Gislaine *et al.*, 2012).

Z. chalybeum has quinoline alkaloids (Kato *et al.*, 1996) and organic extract has flavonoids, alkaloids, saponins and sesquiterpene lactones (Musila *et al.*, 2013).



Figure 4: *Zanthoxylum chalybeum* (Rutaceae) tree
(Photograph: Magrate Kaigongi in Tharaka-Nithi County)

The plants used in this study were selected on the basis of their ethnopharmacological use by the Msambweni community. The preparation for treatment of various ailments in the traditional medicine, chemical constituent and the reported ethnomedical uses for each plant are tabulated below (Table 1).

Table 1: Selected plant species from Msambweni district used in traditional medicine

Family/Species/(Voucher specimen number)	Habit	Part used	Treatment preparation	chemical constituents	Relevant reported ethnomedical uses
<i>Adansonia digitata</i> L. (Bombacaceae) JN01	Tree	Leaf	Decoction	Proanthocyanidins, terpenoids, flavonoids, and sterols have been isolated from <i>A. digitata</i> (VanStaden <i>et al.</i> , 2012).	Leaf decoction is used as antimalarial, diaphoretic and prophylactic against fevers (Nguta <i>et al.</i> , 2010)
<i>Grewia trichocarpa</i> Hochst. Ex A. Rich (Tiliaceae) JN02	Shrub	Root	Decoction	No previous reports	Root decoction is used to treat malaria (Nguta <i>et al.</i> , 2011)
<i>Launaea cornuta</i> (Hocht.ex. Oliv. & Hiern.) C. Jeffrey (Compositae) JN03	Herb	Leaf	Decoction	flavonoids, alkaloids and saponins with sesquiterpene lactone (Musila <i>et al.</i> , 2013)	Root and leaf decoction used to treat malaria and typhoid respectively (Nguta <i>et al.</i> , 2010; Kokwaro, 2009)
<i>Zanthoxylum chalybeum</i> . Engl. (Rutaceae) JN04	Tree	Root bark	Decoction	quinoline alkaloids (Kato <i>et al.</i> , 1996) and organic extract has flavonoids, alkaloids, saponins and sesquiterpene lactones (Musila <i>et al.</i> , 2013).	Leaf decoction is rubbed on swelling for the treatment of oedema in kwashiorkor and snake bites, root decoction is drunk for the treatment of chest pain, pneumonia and bark decoction drunk for the treatment of malaria, colds, coughs, dizziness and also chewed for toothache (Kokwaro, 2009) bark of the Kenyan plant was reported active against <i>Bacillus subtilis</i> , <i>Penicillium crustosum</i> and <i>The Saccharomyces cerevisiae</i> (Taniguchi <i>et al.</i> , 1978)

2.8. Toxicity Studies

Many substances we handle daily are toxic. Different organisms respond differently to the same toxin. Lethal Dose 50 is a term which is used to describe acute toxicity. It refers to lethal

dose which is acutely lethal to 50% of all the organisms in a given experiment. LD₅₀ is expressed in mg/kg or ppm (parts per million) (Shankar *et al.*, 1980). To meet the criteria of efficacy, safety and quality control like synthetic drug products, the pharmacological, toxicological and phytochemical profiles of the plant extracts have to be scientifically evaluated (Wagner, 1997).

2.8.1. Significance of toxicity studies

The use of plant extracts in treatment of diseases without any standard dosage accompanied with lack of adequate scientific studies has raised concern on their toxicity. Toxicity studies help in assessing the right dosage to be administered without causing health risks in the organisms (Ashafa *et al.*, 2012). The lack of precise dosage in traditional medicine is not peculiar as it also occurs in modern medicine due to imprecise diagnosis (Kokwaro, 2009). The toxicity testing using Brine shrimp is cheap, easy, quick and only small amount of the sample to be tested is required. Thus it is an appropriate bioassay of determining the LC₅₀ of various extracts (Apu *et al.*, 2010).

Simple bioassays for determination of bioactivities of crude plant extracts are important for the purpose of standardization or quality control of plant botanicals. Due to this Brine shrimp lethality test is a “top bench” procedure useful in natural product chemistry (Jerry and Lingling, 1998). Using Brine shrimp in toxicity studies is economical, easy, uses small amount of test material, inexpensive, simple and requires little chemical training. This test is a convenient probe for preliminary assessment of toxicity of natural products from plant biodiversity, heavy metals and pesticides (Aseer *et al.*, 2009).

2.8.2. Brine shrimp lethality assay

The shrimp lethality assay was proposed by Michael (Krishanaraju *et al.*, 2005) and later developed (Vanhaecke *et al.*, 1981; Sleet and Brendel 1983). It is based on the ability of pharmaceutical compounds to kill laboratory-cultured *Artemia* nauplii brine shrimp. The brine shrimp lethality bioassay (BSLB) has been used extensively in the primary screening of the crude extracts as well as the isolated compounds to evaluate the toxicity towards brine shrimps, which could also provide an indication of possible cytotoxic properties of the test materials (Meyer *et al.*, 1982).

Since BLSB is the simple method useful for screening large number of extracts in the drug discovery process. The method allows the use of smaller quantity of the extracts and permits larger number of samples and dilutions within shorter time than using the original test vials (Sam, 1993). Furthermore it has been established that the cytotoxic compounds generally exhibit significant activity in the BSLB, and this assay can be recommended as a guide for the detection of antitumour and pesticidal compounds because of its simplicity and low cost (Mazid *et al.*, 2008). This bioassay has also a good correlation with the human solid tumour cell lines. The inhibitory effect of the extract might be due to the toxic compounds present in the active fraction that possess ovicidal and larvicidal properties. The metabolites either affected the embryonic development or slay the eggs. Therefore the cytotoxic effects of the plant extracts enunciate that it can be selected for further cell line assay because there is a correlation between cytotoxicity and activity against the brine shrimp nauplii using extracts (Manilal *et al.*, 2009).

2.9. Clinical and Laboratory Standards Institute protocol on evaluation of antimicrobial activity

The test plates are inoculated with microorganism by optimally dipping a sterile cotton swab into a suspension adjusted to turbidity of 0.5 McFarland scale. The dried surface of agar plate is inoculated by streaking the swab over the agar surface. This procedure is repeated twice by rotating the plate approximately 60° each time to ensure even distribution of inoculums. The discs are applied on the inoculated agar by gently pressing them down to ensure complete contact with the agar. Whether the discs are applied individually or with a dispensing apparatus, they must be distributed evenly so that they are no closer than 24mm from center to center. Ordinarily, no more than 12 discs in one plate. It is also important to pay attention to how close the discs are to the edge of the plate. The plates should be inverted and placed in an incubator set to 37°c for 24hours and 30°c for 48 hours for fungi (CLSI, 2006).

2.10. Interpretation of antimicrobial results

The “susceptible” category implies that isolates are inhibited by the usually achievable concentrations of antimicrobial agent when the recommended dosage is used for the site of infection. The “intermediate” category includes isolates with antimicrobial agent MICs that approach usually attainable blood and tissue levels and for which response rates may be lower than for susceptible isolates. The intermediate category implies clinical efficacy in body sites where the drugs are physiologically concentrated (e.g., quinolones and β-lactams in urine) or when a higher than normal dosage of a drug can be used (e.g., β-lactams). This category also includes a buffer zone, which should prevent small, uncontrolled, technical factors from causing major discrepancies in interpretations, especially for drugs with narrow pharmacotoxicity margins. The “resistant” category implies that isolates are not inhibited by the usually achievable concentrations of the agent with normal dosage schedules, and/or that

demonstrate zone diameters that fall in the range where specific microbial resistance mechanisms (e.g., beta-lactamases) are likely, and clinical efficacy of the agent against the isolate has not been reliably shown in treatment studies. A microorganism for example is said to be resistant against 40µg/ml of gentamicin if the zone of inhibition exhibited is ≤ 12 mm, intermediate at 13-14mm and susceptible at ≥ 15 mm. The commercially manufactured antibiotics and isolated pure compounds exhibit larger zones of inhibition at low concentrations compared to crude extracts (CLSI, 2006). According to Kitonde *et al.* (2013), the lowest MICs ≤ 20 mg/ml indicate highest activity, while 31.25-62.5 mg/ml indicate moderate activity, and ≥ 125 mg/ml indicate low activity

A crude extract with LC₅₀ values ranging between 0-500 µg/ml is said to be highly toxic while that with LC₅₀ values between 500-1000 µg/ml is termed as moderately toxic. An extract with LC₅₀ values above 1000 µg/ml is non toxic (Nguta *et al.*, 2011)

2.11. Problem statement

Microbial infections have caused a big burden of diseases. An alarming increase in microbial resistance against conventional antibiotic treatment has developed over the last forty years in both medical and livestock sectors. This alarming increase in the development of resistance to the antibiotics in the clinical practice has led to the incidence of new and re-emergence of infectious diseases making them expensive and difficult to treat. Plants have great potential as antimicrobial agents due to the presence of a wide range of bioactive compounds such as alkaloids, flavonoids, saponins and sesquiterpene lactones. However, since many medicinal plants are unevaluated, it is possible that the active compounds of many plants go undetected while some plants used therapeutically result in poisoning. Evaluation of chemical activity and toxicity of plants extracts is important in determining safe and efficient dosages. This study was carried out to determine the antimicrobial activity, phytochemical and toxicity analysis of;

Zanthoxylum chalybeum, *Adansonia digitata*, *Launaea cornuta* and *Grewia trichocarpa* crude extracts.

2.12. Justification

In Kenya antibiotic resistance has led to significant morbidity and mortality which is a drawback to economic development. Antibiotics are expensive to poor communities in developing countries and most microbes such as methicilin resistant *S. aureus* have become resistant to almost all available drugs. Research into new alternative for such microbes is warranted. The World Health Organisation (WHO, 2001) advocates for traditional medicine as a safer remedy for ailments of microbial and non-microbial origin. Most modern medicines were discovered through study of plants which were used traditionally to treat specific illnesses. In addition, very few medicinal plants have been analyzed chemically and their bioactive constituents are yet to be validated. As a result, knowledge from traditional medicine can be very essential in the development of cheap and effective antibiotics.

2.13. Objectives

2.13.1. Main Objective

To determine *in vitro* antimicrobial activity, toxicity and phytochemical composition of crude extracts from *Zanthoxylum chalybeum* , *Adansonia digitata*, *Launaea cornuta* and *Grewia trichocarpa*.

2.13.2. Specific objectives

1. To evaluate the antimicrobial potential of selected aqueous and organic crude plant extracts
2. To determine the acute toxicity of the crude extracts on Brine shrimp larvae
3. To characterise the major phytochemical compounds in the crude plant extracts.

2.13. 3. Null Hypothesis

Crude extracts of *Zanthoxylum chalybeum*, *Adansonia digitata*, *Launaea cornuta* and *Grewia trichocarpa* will have activity against *in vitro* growth of methicilin resistant *Staphylococcus aureus*, *Bacillus cereus*, *Escherichia coli*, *Pseudomonas aeruginosa* and *Candida albicans* and will be toxic to *Artemia salina* larvae

CHAPTER THREE: MATERIALS AND METHODS

3.1. Collection of plant materials

The various plants parts depending on the ethnobotanical knowledge on specific plants were collected from Msambweni District in Kenya by Dr. Joseph Nguta of University of Nairobi. The selection of the four plants was done on the basis of their wide usage in treatment of not only microbial infections but also malaria (Nguta et al., 2011) by the Msambweni community. Plants were identified by a taxonomist from the University of Nairobi and voucher specimens deposited in the University of Nairobi herbarium. The plant parts were air dried at room temperature, chopped into small pieces and ground into powder.

Table 2: Plants Collected from Msambweni District, Kenya

Voucher specimen number	Plant species	Family	Plant Part Collected
JN01	<i>Adansonia digitata</i> L	Bombacaceae	Stem Bark
JN02	<i>Grewia trichocarpa</i> Hochst. Ex A. Rich	Tiliaceae	Roots
JN03	<i>Launaea-cornuta</i> (Hocht.ex.Oliv.&Hiern.) C.Jeffrey	Compositae	Leaves
JN04	<i>Zanthoxylum chalybeum</i> .Engl.	Rutaceae	Root bark

3.2. Preparation of Crude Extracts

The ground materials were extracted by cold maceration method. 50gm of ground plant material in 500mls of distilled water to obtain aqueous extracts and 50gms in 500mls of chloroform-Methanol mixture (1:1) for 48 hours to obtain organic extracts. The aqueous extracts were filtered and the filtrate kept in a deep freezer then lyophilized (freeze dried) resulting to a dry powder. Organic extracts were filtered and concentrated with a rotary

evaporator to dry powder (Nguta *et al.*, 2011). The resulting dry powders were used in antimicrobial, phytochemical screening and toxicity evaluation.

3.3. Microorganism Suspension Standardization

The microorganisms used in the study were obtained from the Department of Public Health, Pharmacology and Toxicology, University of Nairobi and the Centre for Microbiology Research (CMR), Kenya Medical Research Institute (KEMRI). The fungal suspension was standardized according to the Clinical and Laboratory Standards Institute (CLSI, 2008). The yeast was grown on Sabouraud agar for 72 hours and standardized with sterile saline to turbidity equivalent to 0.5 McFarland standards which is approximately $1-5 \times 10^6$ CFU/ml. Bacteria were grown in Muller–Hinton agar for 24 hours and standardized with sterile saline to turbidity equivalent to 0.5 McFarland scale approximately $1-2 \times 10^8$ CFU/ml (CLSI, 2009) and stored at 4°C. The antibacterial and antifungal activity was determined using disc diffusion test and broth dilution method (Kalayou *et al.*, 2012).

Table 3: List of microbes tested in the study

Name of microbe	Microbe and type used	Details of strain used	Source
<i>Bacillus cereus</i>	Gram positive bacteria	ATCC 11778	DPHPT, UoN
Methicilin Resistant <i>S. aureus</i> (MRSA)	Gram positive bacteria	Untyped pure clinical isolate collected from CMR- KEMRI	KEMRI
<i>Pseudomonas aeruginosa</i>	Gram negative bacteria	ATCC 27853	DPHPT, UoN
<i>Escherichia coli</i>	Gram negative bacteria	ATCC 25922	DPHPT, UoN
<i>Candida albicans</i>	Fungal yeast	ATCC 10231	DPHPT, UoN

The microbes were selected on the basis of their availability.

3.4. Disc Diffusion Test

The disc diffusion method followed by the Clinical and Laboratory Standards Institute protocol was used to evaluate antimicrobial activities (Kaloyou *et al.*, 2012). To determine susceptibility, four concentrations (200 mg/ml, 100 mg/ml, 50 mg/ml and 25 mg/ml of crude extracts) were prepared in dimethylsulfoxide (DMSO) and water for organic and aqueous extracts respectively by dissolving 2g of each extracts in 10mls and serially diluted to make 100mg/ml, 50mg/ml and 25mg/ml. Sterile antibiotic assay discs (Whatman, 6 mm) were impregnated with 100 μ L of the reconstituted extract and dried completely under sterile conditions in a laminar flow. Each disc was gently pressed down to ensure complete contact with the agar inoculated with 1ml of test microorganisms (bacteria and fungi). Extracts were tested in triplicate. DMSO and water saturated assay discs were used as negative control. Gentamicin and Amphotericin B were used as positive control for bacteria and fungi respectively. The plates were incubated at 37 °C for 18-24 h for bacteria and 30°C for 72 hrs for fungi. Inhibition zones were recorded as the diameter of growth free zones.

3.5. Determination of minimum inhibitory concentration (MIC)

The extracts that showed antibacterial activity against the selected pathogens were selected for further tests to calculate their MIC by dilution method. This test was performed in sterile 96-well microplates. Microdilution was performed in 96-well microtiter plates with U-shaped wells where the cultures were diluted in Müeller-Hinton broth at a density adjusted to 0.5 McFarland turbidity. Controls with 0.5 ml of only culture medium or others with plant extracts were used in the tests. The wells were filled with 100 μ l of sterile H₂O and 100 μ l of the plant extracts were added to the wells by serial two fold dilution from the suspension of plant extract stock solution. Each well was inoculated with 100 μ l of bacterial and fungal suspensions. The plates were covered, placed in plastic bags to create anaerobic conditions and incubated at

37°C for 24 hrs for bacteria and 30°C for 72hrs for fungi. This was followed by inoculating the cultures on the agar. In this study, the MIC was the lowest concentration of plant extracts that exhibited no growth of the microorganism (Kalayou *et al.*, 2012)

3.6. Acute toxicity testing

The toxicity of the extracts (Both the aqueous and organic) was determined using Brine shrimp-Leech (*Artemia salina L.*) eggs (Wanyoike *et al.*, 2004). A hatching tank measuring 14cm by 5cm and 9cm was used. The tank was divided into two unequal compartments by a plastic divider with holes of diameter of 2mm. The tank was filled to the brim using sea water. The sea water was prepared by dissolving 30g of commercial sea salt in a one liter of distilled water. A spoonful of *Artemia salina* eggs and a gram of yeast to act as food for the nauplii were added into the larger compartment of the hatching tank containing the sea water. This was followed by covering the larger compartment to enhance hatching of the eggs by preventing light penetration. This was incubated at room temperature (22-29°C) for 48hrs to allow hatching of the eggs.

Various concentrations of the crude extract in sea water were used: 10,100 and 1000µg/ml. A stock solution for both the organic and aqueous crude extract of 10,000µg/ml was prepared from which serial dilutions were done to make the three concentrations used in the experiment. The aqueous extracts stock solution of 10,000µg/ml was prepared by dissolving 0.1g of the crude extract in 10 mls of distilled water and the chloroform: MeOH extracts, 0.1g of the sample was first dissolved in dimethylsulphoxide (DMSO) then diluted further using artificial water to 10 mls to make stock solution of 10,000µg/ml. The stock solutions from the crude extracts were filtered using 0.22 µm micro filters under lamina flow hood (Wanyoike *et al.*, 2004). The solutions were then put into the pre-marked vials and 10 live brine shrimp nauplii were added in 5 ml simulated sea water (Apu *et al.*, 2010). Each experiment was replicated

three times. A negative control was prepared with artificial sea water and DMSO only for organic extract and artificial sea water only in the case of aqueous extract (Wanyoike *et al.*, 2004). Survivors were counted after 24 hrs using a dissecting microscope. The mortality endpoint of this bioassay was defined as the absence of controlled forward motion during 30 s of observation (Apu *et al.*, 2010).

3.7. Phytochemical Screening of Plants Extracts Using Thin Layer Chromatography (TLC)

The crude plant extracts were screened for flavonoids, alkaloids, sesquiterpene lactones and saponins. This was done by dissolving the organic and aqueous extracts in chloroform and methanol (1:1). Using capillary tubes, spots of the dissolved extracts were made on the base line of the aluminium TLC plates. The plates were put in a chamber containing appropriate chromatographic solvent system (mobile phases) specific for the determination of the presence of a given class of phytochemical constituents (Harbourne, 2002)

Table 4: Procedure followed in screening for different chemical compounds

Class of Secondary metabolites	Solvent system for developing TLC plates	Detection
Alkaloids	Dichloromethane: Methanol (85:15) (Harborne, 2002)	Dragendorff's reagent was sprayed on the developed plates. Formation of orange colors indicated the presence of alkaloids. Spraying the plates with sodium nitrate made the orange colors more intense (Harborne, 2002)
Flavonoids	n-hexane: ethyl acetate: acetic acid (6:3:1) (Waksmundzka <i>et al.</i> , 2008)	At 254 nm, flavonoids appeared as dark blue zones on a yellow background on the developed plates. The intensity of the yellow color in the background increased when ammonia was sprayed on the plates. At 365 nm flavonoids fluoresced yellow, blue or green (Waksmundzka <i>et al.</i> 2008).
Saponins	Dichloromethane:Ethyl acetate (9:1) according to Karem <i>et al.</i> (2005)	Detection of saponins occurred when plates were sprayed with a mixture of ethanol and H ₂ SO ₄ (9:1) and then heated at 110 ⁰ C for 10 minutes. Saponins appeared as black spots (Karem <i>et al.</i> , 2005)
Sesquiterpene lactones	n- hexane: ethyl acetate (9:1) (Waksmundzka <i>et al.</i> , 2008).	Detection was done by placing the developed plates in a chamber containing iodine crystals to observe brown spots (Waksmundzka <i>et al.</i> , 2008). Sesquiterpene lactones were also detected as brown, yellow spots when plates were sprayed with concentrated H ₂ SO ₄ and heated for 5 minutes at 100-110 ⁰ C (Harborne, 2002).

3.8. Data analysis

Statistical analysis of antimicrobial activity was done using statistical program for social sciences (statistical analysis software). Using the software, ANOVA was used to determine whether there were significant differences in the mean diameter of inhibition zones in various concentrations. Once the means were found to be different from each other, Dunnett test was then used for multiple comparisons of inhibition to determine whether inhibition arising from the various treatments were different from the inhibition induced by the positive controls. The significance level used in the analysis was 0.05 (Alpha Level \leq 0.05).

In toxicity testing, LD_{50} was determined by Finney's probit analysis. Probit analysis is a specialized regression model of binomial response variable and it has been used to analyse dose-response experiments in a variety of fields. It involves conversion of concentrations into logarithms and the corresponding percentage mortalities into probits. The logarithms of the concentrations are then plotted against the probits to give a straight line graph. This regression line is then used in the determination of LD_{50} (Finney, 1971).

CHAPTER FOUR: RESULTS

4.1 Extraction yields

The resulting dry powders were weighed and expressed as percentages. The ground powders of organic extracts of *A. digitata*, *G. trichocarpa*, *L. cornuta* and *Z. chalybeum* yielded 1.1 %, 1.7 %, 2.8 % and 6.3 % respectively. The ground powders of aqueous extracts of yielded *A. digitata*, *G. trichocarpa*, *L. cornuta* and *Z. chalybeum* yielded 1.3 %, 1.6 %, 4.1 % and 3.2 % respectively. *Z. chalybeum* organic extract yielded the highest recovery value of 6.3 % while *A. digitata* organic extract had the lowest yield of 1.1 %. (Table 5).

Table 5: Percentage yield of plant extracts using aqueous and organic extraction methods

Plant species	Part used	Extraction type	% yield to weight of dry powered plant
<i>Adansonia digitata</i>	Stem Bark	Organic	1.1
<i>Adansonia digitata</i>	Stem Bark	Aqueous	1.3
<i>Grewia trichocarpa</i>	Root	Organic	1.7
<i>Grewia trichocarpa</i>	Root	Aqueous	1.6
<i>Launnaea cornuta</i>	Leaves	organic	2.8
<i>Launnaea cornuta</i>	Leaves	Aqueous	4.1
<i>Zanthoxyum chalybeum</i>	Root bark	Organic	6.3
<i>Zanthoxyum chalybeum</i>	Root bark	Aqueous	3.2

4.2. Antimicrobial activity

4.2.1. Disc Diffusion Technique


4.2.1.1. Aqueous extracts against *B. cereus*

The aqueous crude extracts of *A. digitata* and *Z. chalybeum* showed antimicrobial activities against *B. cereus* at 200 mg/ml and 100 mg/ml. *A. digitata* showed inhibition zones of 8.67 mm and 7.5 mm at 200 and 100 mg/ml respectively while that of *Z. chalybeum* was 10.5 mm and 7.67 mm at both 200 and 100 mg/ml respectively. The aqueous crude extracts of *G. trichocarpa* and *L. cornuta* did not show any antimicrobial activity against *B. cereus*

4.2.1.2. Organic extracts against *B. cereus*

Organic crude extract of *Z. chalybeum* showed the highest inhibition against *B. cereus* at 200 and 100 mg/ml concentrations with 13.87 mm and 12.167 mm respectively. The growth inhibition (mm) of *Z. chalybeum* organic extract at 200 mg/ml against *B. cereus* did not show much difference from that of the positive control (Gentamicin 0.04 mg/ml), $P \geq 0.05$ as shown in figure 5.

Inhibition zone exhibited by *Z. chalybeum* (200mg/ml) against *B. cereus*



Inhibition zone exhibited by the gentamcin (0.04 mg/ml) against *B. cereus*




Figure 5: Inhibition zones caused by *Z. chalybeum* at 200 mg/ml and positive control against *B. cereus*

A. digitata organic extract also showed antimicrobial activity against *B. cereus* at 200 mg/ml and 100 mg/ml with inhibition zones of 8.67 mm and 7.5 mm respectively. *G. trichocarpa* showed antimicrobial activity at 200, 100 and 50 mg/ml with inhibition zones of 11.67, 9.67 and 7.17 mm respectively (Figure 6).




Figure 6: Growth inhibition of *B. cereus* by different concentrations of extracts from the four test plants

4.2.1.3. Aqueous extracts against *P. aeruginosa*

In aqueous extracts of all the tested plants, only *A. digitata* inhibited *P. aeruginosa*. *A. digitata* showed antimicrobial activity 200 mg/ml, 100 mg/ml and 50 mg/ml with inhibition zones of 9.17 mm, 8 mm and 7.5 mm respectively.

4.2.1.4. Organic extracts against *P. aeruginosa*

The organic extracts of *G. trichocarpa* showed the highest levels of inhibition in all concentrations with 10.67, 9.67, 8.83 and 7.33 mm respectively. *Z. chalybeum* showed inhibition zones of 10.5 mm, 9.17 mm, 8 mm and 7.67 mm at 200 mg/ml, 100 mg/ml, 50 mg/ml and 25 mg/ml respectively. *A. digitata* and *L. cornuta* organic extracts showed

antimicrobial activities at 200, 100 and 50 mg/ml with *A. digitata* showing inhibition zones of 9.17 mm, 9 mm and 8.5 mm respectively while *L. cornuta* showed inhibition zones of 10.5 mm, 9.17 mm and 8 mm respectively (figure 7)




Figure 7: Growth inhibition of *P. aeruginosa* by different concentrations of extracts from the four test plants

4.2.1.5. Aqueous Extracts Against Methicillin Resistant *S. aureus* (MRSA)

In the case of MRSA, none of the aqueous extracts showed any inhibition except *A. digitata* which showed inhibition zones of 9.33 mm and 8.17 mm at 200 mg/ml and 100 mg/ml respectively.

4.2.1.6. Organic Extracts against MRSA

All organic extracts showed some inhibition against MRSA. *A. digitata* organic extract at 200 mg/ml and 100 mg/ml showed highest inhibition zones of 14.33 mm and 12 mm respectively. *A. digitata* organic extract zone of inhibition (mm) at 200 mg/ml against MRSA was not much different from that of the positive control (Gentamicin, 0.04 mg/ml), $P \geq 0.05$. The organic extract of *G. trichocarpa* was the second in terms of activity at 200 mg/ml with 10.33 mm followed by the organic extract of *Z. chalybeum* at 200 mg/ml with 9.33 mm. All the organic extracts showed inhibition at 50 mg/ml while *A. digitata* showed inhibition of 8.33 mm at the lowest concentration 25 mg/ml (Figure 8).




Figure 8: Growth inhibition of MRSA by different concentrations of extracts from the four test plants

4.2.1.7. Aqueous extracts against *C. albicans*

The aqueous extracts of the four plants did not inhibit the growth of *C. albicans* at any of the concentration used.

4.2.1.8. Organic extracts against *C. albicans*

All the organic extracts except for *L. cornuta* showed inhibition against *C. albicans* at 200 and 100 mg/ml. *A. digitata* registered the inhibition of 9.7 mm and 8.3 mm at 200 mg/ml and 100 mg/ml respectively. The inhibition zones caused by *A. digitata* organic extract at 200 mg/ml and Amphotericin B against *C. albicans* is shown in figure 9.




Figure 9: Inhibition zones showing antifungal activity of *A. digitata* at 200 mg/ml and positive control against *C. albicans*

Z. chalybeum at 200 mg/ml and 100 mg/ml showed inhibition of 8.7 mm and 7.5 mm respectively while *G. trichocarpa* showed inhibition of 8.5 mm and 7.83 mm at 200 mg/ml and 100 mg/ml respectively (Figure 10).




Figure 10: Growth inhibition of *C. albicans* by different concentrations of extracts from the four test plants

4.2.2. Minimum Inhibitory Concentration (MIC) Results

Extracts which at 200 mg/ml did not inhibit the microorganism was assumed to have MIC value greater than 200 mg/ml.

4.2.2.1. Organic Extracts

Minimum inhibitory concentration (MIC) values of the extracts tested against methicillin resistant *S. aureus* (MRSA) shows *A. digitata* organic extract had the highest activity with the lowest MIC of 20 mg/ml. *Z. chalybeum* and *G. trichocarpa* organic extracts had the highest activity against *P. aeruginosa* with MIC of 20 mg/ml. *Z. chalybeum* organic extract exhibited the highest activity against *B. cereus* at 200 mg/ml with an inhibition zone of 13.17 mm, however *G. trichocarpa* organic extract showed the lowest MIC value against *B. cereus*. *Z. chalybeum* organic extract exhibited highest activity against *C. albicans* with a MIC of 40 mg/ml (Table 6).

Table 6: Minimum Inhibitory Concentration (MIC) of Organic Crude Extracts

PLANT NAME	MIC (mg/ml)				
	<i>C. albicans</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>B. cereus</i>	MRSA
<i>Adansonia digitata</i>	70	> 200	30	90	20
<i>Zanthoxylum chalybeum</i>	40	> 200	20	60	40
<i>Launnaea cornuta</i>	> 200	> 200	40	> 200	40
<i>Grewia trichocarpa</i>	80	> 200	20	40	40

4.2.2.2. Aqueous Extracts

Some aqueous extracts did not show any antimicrobial activity against most of the microbes at any concentration tested and therefore their MIC values were assume to be more than 200 mg/ml. *A digitata* recorded MIC values of 40 mg/ml against *P. aeruginosa*, 90 mg/ml against *B. cereus* and 80 mg/ml against MRSA while *Z. chaybeum* had MIC value of 90 mg/ml against *B. cereus* (Table 7)

Table 7: Minimum Inhibitory Concentration (MIC) of Aqueous Crude Extracts

PLANT NAME	MIC (mg/ml)				
	<i>C. abicans</i>	<i>E. coli</i>	<i>P. aeruginosa</i>	<i>B. cereus</i>	MRSA
<i>Adansonia digitata</i>	> 200	> 200	40	90	80
<i>Zanthoxylym chalybeum</i>	> 200	> 200	> 200	90	> 200
<i>Launnaea cornuta</i>	> 200	> 200	> 200	> 200	> 200
<i>Grewia trichocarpa</i>	> 200	> 200	> 200	> 200	> 200

4.3. Brine-Shrimp lethality test

4.3.1. Aqueous Crude Extracts

A. digitata aqueous crude extract had the highest LC₅₀ value of 3988 µg/ml followed by *G. trichocarpa* with 1488 µg/ml while *Z. chalybeum* exhibited the least LC₅₀ value of 293.5 as shown in table 8

Table 8: LC₅₀ values for various tested aqueous crude extract on Brine-shrimp larvae

Plant species	Average mortality at various concentrations				LC ₅₀ µg/ml
	1,000 µg/ml	100 µg/ml	10 µg/ml	0 µg/ml	
<i>Adansonia digitata</i>	1.33	0.33	0	0	3988
<i>Zanthoxylym chalybeum</i>	8	1	0	0	293.5
<i>Launnaea cornuta</i>	6.33	1.67	0.33	0	554.4
<i>Grewia trichocarpa</i>	0.33	0	0	0	1488

4.3.2. Organic Extracts

A. digitata organic crude extract showed the highest LC₅₀ value of 5626 followed by *G. trichocarpa* with 5602. *L. cornuta* showed LC₅₀ value of 141.3 while *Z. chalybeum* recorded the lowest LC₅₀ value of 39.2 (table 9)

Table 9: LC₅₀ values for various tested organic crude extract on Brine-shrimp larvae

Plant species	Average mortality at various concentrations				LC ₅₀
	1,000 µg/ml	100 µg/ml	10 µg/ml	0 µg/ml	µg/ml
<i>Adansonia digitata</i>	2.67	1	0	0	5626
<i>Zanthoxylum chalybeum</i>	10	5.67	3	0	39.2
<i>Launnaea cornuta</i>	8.33	4	1.33	0	141.3
<i>Grewia trichocarpa</i>	2.33	0	0	0	5602

4.4. Phytochemical screening of secondary compounds




Figure 11: A photograph showing sesquiterpene lactones results of the tested extracts on a TLC plate

A- *A. digitata* organic, B- *L. cornuta* organic, C- *Z. chalybeum* organic, D- *G. trichocarpa* organic, E- *A. digitata* aqueous, F- *L. cornuta* aqueous, G- *Z. chalybeum* aqueous and H- *G. trichocarpa* aqueous

4.4.1. Aqueous Crude Extracts

Alkaloids and flavonoids were present in all the aqueous extracts while saponins were only found in *L. cornuta* only. Sesquiterpene lactones were absent in all the aqueous extracts (table10)

Table 10: Presence of alkaloids, flavonoids, sesquiterpene- lactones and saponins in aqueous extracts

Extract	alkaloids	flavonoids	Sesquiterpene lactones	saponins
<i>Adansonia digitata</i>	+	+	-	-
<i>Zanthoxylym chalybeum</i>	++	+	-	-
<i>Launnaea cornuta</i>	+	+	-	+
<i>Grewia trichocarpa</i>	+	+	-	-

-...absence

+...presence, the number of (+) shows comparative amounts depending on degree of coloration

(b).Organic Crude Extracts

Alkaloids, flavonoids, Sesquiterpene lactones and saponins were present in all the organic extracts (table 11)

Table 11: Presence of alkaloids, flavonoids, sesquiterpene- lactones and saponins in organic extracts

Extract	alkaloids	flavonoids	Sesquiterpene lactones	saponins
<i>Adansonia digitata</i>	++++	+	+	++
<i>Zanthoxylum chalybeum</i>	+++	+++	+	+++
<i>Launnaea cornuta</i>	+	++	+++	++
<i>Grewia trichocarpa</i>	++	+	++	+

-...absence

+...presence, the number of (+) shows comparative amounts depending on degree of coloration

CHAPTER FIVE: DISCUSSION

The current study was designed to investigate the antimicrobial activity, toxicity and phytochemical screening of four medicinal plants used in Msambweni District, Kenya. Due to the side effects of the current drugs the resistance that pathogenic microorganisms build against antibiotics, much attention has led to study of biologically active compounds isolated from plant species used in herbal medicine (Essawi and Srour, 2000). The results of different studies provide evidence that medicinal plants might indeed be potential sources of new antibacterial agents even against some antibiotic-resistant strains (Kone *et al.*, 2004).

From this study, it is clear that the chloroform: methanol (1:1) solvent extracts of all the plants tested were more potent than their corresponding aqueous extracts against all the tested microbes. *A. digitata* organic extract was active against all the microbes except the *E. coli*. The highest activity of *A. digitata* recorded was against MRSA followed by *P. aeruginosa* and *B. cereus*. The high antimicrobial activity of organic extract of *A. digitata* stem bark against MRSA may be used as yardstick to identify novel drugs with the potential to combat bacterial strains resistant to known antibiotics, shorten the length of treatment, increase patient compliance, and avoid overdose which may lead to toxicity or side effects to patients. These results are in agreement with VanStaden *et al.*, (2011) who reported that *A. digitata* has considerable antimicrobial activity against *Bacillus subtilis*, *S. aureus* and *C. albicans*. This signifies that, *A. digitata* extract has potential novel antimicrobial compounds for management of diseases caused by the above microbes.

Launnaea cornuta aqueous extract did not show any inhibition against the tested microbes at any concentration. However, the organic extract showed moderate activity against *P. aeruginosa* and MRSA and failed to inhibit *B. cereus*, *E. coli* and *C. albicans*. This is in

agreement with findings from Nagata *et al.* (2011) who reported that leaves and roots of *L. cornuta* are used as antimicrobial by the people of Suba District-Kenya particularly those living with HIV/AIDS to manage opportunistic diseases. However, the low activity exhibited by *L. cornuta* in this study may be attributed to the difference in the dose used in this study, the method of extraction of medicinal plants, the method of antimicrobial study, the genetic variation of plant, age of the plant or the environment.

Zanthoxylum chalybeum showed antimicrobial activity with the highest activity being observed against *B. cereus* whereby at 200 mg/ml, there was no much difference in terms of inhibition zones (mm) with that of standard antibiotic (Gentamicin, 0.04 mg/ml) ($P \geq 0.05$). *Z. chalybeum* organic extract also showed high activity against *P. aeruginosa* and moderate activity against MRSA and *C. albicans* but failed to inhibit the growth of *E. coli* at all tested concentrations. These results are in agreement with studies by Adesina's (2005) findings that the ethanol extract of *Z. chalybeum* has antibacterial activity against *B. cereus* and *P. aeruginosa* and antifungal activity against *C. albicans*.

Grewia trichocarpa organic extract (CHCl₃: MeOH) showed considerable antimicrobial activity against four of the five tested microbes; MRSA, *B. cereus*, *P. aeruginosa* and *C. albicans* and failed to show any inhibition against *E. coli*. The aqueous extract of *G. trichocarpa* did not inhibit any of the tested microbes. The organic extract showed high activity against *P. aeruginosa* and moderate activity against *C. albicans*, *B. cereus* and MRSA. This is the first report on antimicrobial activity of *G. trichocarpa*.

Differential sensitivity of microbes to the extracts may be explained by the cell wall composition of the gram positive and gram-negative bacteria. The cell wall of gram negative

bacteria contains the outer membrane and lipid bilayer embedded with proteins and porins. This determines and allows different molecules or ions either into or out of the cell and thus the outer membrane serves as a barrier to the passage of many molecules and hence less sensitive to many extracts (Kalayou *et al.*, 2012). This supports why all the extracts did not inhibit the growth of *E. coli* which was one of the gram negative bacteria. The gram positive bacterium has a relatively thick layer of peptidoglycan sheets of interconnected glycan chains made up of polymer which is fully permeable to many substances, and thus sensitive to most extracts, Nester *et al.*, (2004). This may explain why the gram positive bacteria MRSA and *B. cereus* in this study were susceptible to most extracts.

Low activity of the crude extracts against *C. albicans* (yeast fungus) compared to the bacteria may also be due to differences in cell wall composition of the organisms. Yeast fungus cell wall contains polysaccharides and proteins (Nester *et al.*, 2004). The protein expression on *C. albicans*, function as selective transport system used to expel wastes and compounds that are otherwise deleterious to the cell; a function called efflux which is medically important in that, it allows micro-organisms to oust antimicrobial medications that are made to destroy them and therefore render them resistance, Nester *et al.*, (2004). This explanation could be one of the reasons why *C. albicans* was resistant to the tested crude extracts. The observation reported in this study is in agreement with the findings of Masakazu *et al.*, (2010), who did research on antifungal drug resistance of *Candida* spp and reported that expression of drug efflux pumps were responsible for the resistance.

The organic extracts were more active than aqueous extracts. Noting that the traditional herbal remedy preparation from these plants is by use of water, it is a paradox that the aqueous extracts were inactive in this study. This may be due to the absence or insufficient and

ineffective concentration of the antimicrobial constituents in the aqueous extracts. These findings are in agreement with the studies of Koua *et al.*, (2011) who reported that inactivity of aqueous extracts of *Striga hermonthica* may be due to absence or insufficient and effective concentration of the antimicrobial agents of *S. hermonthica* extracts. It is well known that patients using the traditional herbs take in large amounts of the concoctions and hence may eventually consume sufficient amounts of the curative drugs to elicit healing.

The study on toxicity of *A. digitata* against Brine-shrimp confirmed that both the aqueous and organic extract of *A. digitata* were non-toxic since both extracts had $LC_{50} > 1000 \mu\text{g/ml}$, an observation that is in agreement with that of Musila *et al.*, (2013). *A. digitata* has also been shown to be non-toxic on mice from previous studies. For instance, LD_{50} of aqueous extract of the fruit pulp of *A. digitata* on mice was found to be over $8000 \mu\text{g/ml}$ (Ramadan *et al.*, 1994). This corroborates the results obtained in this study on the non-toxicity of stem bark of *A. digitata* on Brine shrimp larvae. Non-toxicity of *A. digitata* explains why most of the plant parts: seeds, fruit pulps and leaves are consumed as food by many communities. The non-toxicity of *A. digitata* stem implies that the plant may be used to inhibit the growth disease-causing bacteria and fungi without eliciting any toxicity on the patients.

Effects of *L. cornuta* on Brine shrimp from this study showed that both the aqueous and organic extracts (CHCl₃: MeOH) were toxic particularly the (CHCl₃: MeOH) ones. Similarly high levels of toxicity of stem bark of *L. cornuta* on Brine shrimp larvae has also been reported by Nguta *et al.* (2011) and Musila *et al.* (2013).

The study on toxicity of *Z. chalybeum* found that both the aqueous and organic (CHCl₃: MeOH) extracts of *Z. chalybeum* were highly toxic against brine shrimp larvae. Nguta *et al.* (2011) while investigating toxicity of aqueous extracts of the leaves, stem bark and root bark of

Zanthoxylum chalybeum on Brine shrimp larvae obtained $LC_{50} < 500 \mu\text{g/ml}$ for the three plant parts. This confirms the results obtained in this study and implies that *Z. chalybeum* may not make safe antimicrobial herbal remedies. This calls upon cautious use of the plant through dose adjustment amongst communities using this plant for preparation of herbal decoctions. Similarly, toxicity of *Z. chalybeum* methanolic root bark extract on human normal fetal lung fibroblast cells has also been reported by Kamuhabwa *et al.* (2000).

Toxicity study showed that both the organic and aqueous extracts of *G. trichocarpa* were non toxic to brine shrimp larvae. This is the first report on the toxicity studies of *G. trichocarpa*.

The organic extract of *G. trichocarpa* contained alkaloids, flavonoids, saponins and sesquiterpene lactones while the aqueous extract contained only alkaloids and flavonoids. There is no available report from the literature on the antimicrobial activity, toxicity and phytoconstituents of *Grewia* species except on *Grewia hexaminta* which is said to contain triterpenoid compounds (Raghunathaiyar, 1996). This could be the first time to report on *G. trichocarpa* phytochemistry.

Thin layer chromatography (TLC) study of *A. digitata* stem bark shows that the organic extract contained alkaloids, flavonoids, saponins and sesquiterpene lactones while the aqueous extract lacked both saponins and sesquiterpene lactones and contained alkaloids and flavonoids only. This is in agreement with Musila *et al.* (2013). In addition, *A. digitata* stem bark contains medicinal compounds which are largely classified under saponins, alkaloids and flavonoids such as lupeol acetate, β -sitosterol, scopoletin, friedelin, betullinic acid and adansonin while the fruit pulp of *A. digitata* is rich in procyanidins (Sidibe & Williams, 2002; Shahat, 2006). Alkaloids, flavonoids and sesquiterpenes have been reported to be potent plant secondary metabolites with broad spectrum of bioactivities (Mazid *et al.*, 2011). The higher antimicrobial

activity of *A. digitata* on MRSA could be attributed to the presence of large amounts alkaloids. Alkaloids are pharmacogenically active basic principles of flowering plants, (Das *et al.*, 2010). The observed activity of these compounds in *A. digitata* is in line with that of Karou *et al.* (2006) who demonstrated that the Indoloquinoline alkaloid causes cell lysis and morphological changes of *S. aureus*.

Secondary metabolites screening in this study found that both aqueous and organic (CHCl₃: MeOH) extracts of *L. cornuta* contained flavonoids, alkaloids, saponins with sesquiterpene lactones lacking in the aqueous extract. These results are in agreement with (Musila *et al.*, 2013). Elsewhere in Ali *et al.* (2003), the genus *Launaea* is characterized for flavonoids, triterpenes, sesquiterpene lactones, coumarins and steroids. Various species of *Launaea* such as, *L. arborescens*, *L. mucronata*, *L. nudicaulis* and *L. capitata* contain various types of flavones such as luteolin, apigenin and flavone glycosides such as apigenin 7-O-glucoside, vitexin, luteolin 7-O-glucoside and luteolin 7-O-rhamnoside besides others (Christian & Vipaporn, 2010).

Organic extract (CHCl₃: MeOH) of *Z. chalybeum* was found to contain alkaloids, flavonoids, saponins and sesquiterpene lactones while the aqueous extract contained only alkaloids and flavonoids. Similar results were obtained by Musila *et al.* (2013). *Z. chalybeum* has been reported to contain alkaloids which have antibacterial and cytotoxic activity (Chrían *et al.*, 2011). *Zanthoxylum* species contains various compounds such as alkaloids, aliphatic and aromatic amides, lignans, coumarins, sesquiterpene lactones and sterols (Cheng *et al.*, 2011; He *et al.*, 2002).

Biological activity is attributed to the presence of various secondary metabolites in plants (Mazid *et al.*, 2011). Not only their presence, but also the quantity of the phytochemical constituents in a given plant extract determines the extent of extracts' bioactivity. In addition, presence of more than one class of secondary metabolites in a given plant extract will also determine the nature and extent of extract's biological activity (Wang *et al.*, 2010). The presence of flavonoids in the crude extracts may explain the antimicrobial activity as flavonoids are known to complex with the extracellular and soluble proteins and bacterial cell wall of bacteria. More flavonoids may also disrupt microbial membrane. The sesquiterpene lactones are known to be active against bacteria, protozoa and viruses through membrane disruption. (Cowan, 1999).

CHAPTER SIX: CONCLUSIONS AND RECOMMENDATIONS

The organic crude extracts of three plants; *A. digitata*, *Z. chalybeum* and *G. trichocarpa* were active against *B. cereus* and *C. albicans* while all the four plants organic crude extracts were active against *P. aeruginosa* and MRSA. None of the tested crude extracts was active against *E. coli*. The current study shows that *Z. chalybeum* and *A. digitata* inhibit the growth of *B. Fcereus* and MRSA respectively with no significant difference ($P \geq 0.05$) with that of standard antibiotic (Gentamicin). The crude extracts of *A. digitata* and *G. trichocarpa* proved to be non toxic on Brine shrimp larvae with $LC_{50} > 1000 \mu\text{g/ml}$ while *Z. chalybeum* and *L. cornuta* were found to be highly toxic with $LC_{50} < 500 \mu\text{g/ml}$.

Flavonoids, alkaloids, saponins and sesquiterpene lactones were present larger amounts in organic extracts of *A. digitata*, *G. trichocarpa*, *L. cornuta* and *Z. chalybeum* compared to their aqueous extracts. Alkaloids and flavonoids were also present in aqueous extracts of *A. digitata*, *G. trichocarpa*, *L. cornuta* and *Z. chalybeum* while saponins were only present in aqueous extracts of *L. cornuta*. Alkaloids were present in high amount while sesquiterpene lactones were present in low amounts in all the extracts.

Laboratory and clinical studies of these plants are required in order to better understand their antimicrobial activities so as allow the scientific community to recommend their uses as an accessible alternative to synthetic antibiotics. The antimicrobial activity, toxicity and phytochemistry of crude extracts *G. trichocarpa* is being reported for the first time. This could be a start point for further drug research of this species on a wide diversity of microbial pathogens. Though there is no harm in exploitation or threat against the use of these potential plant species, they should be protected and probably domesticated for commercial exploitation in large scale

REFERENCES

- Abou-Zeid, A.M., Altalhi, A. D., Abd El-Fattah, R.I. (2008). Fungal Control of Pathogenic Fungi Isolated From Some Wild Plants in Taif Governorate, Saudi Arabia. *Malysian Journal of Microbiology*, 4(1), 30-39.
- Adesina, S.K. (2005). The Nigerian *Zanthoxylum*; chemical and biological values. *African Journal of Traditional and Complimentary medicine*, 2(3) ,282 – 301
- Ahmad, I., & Beg, A. Z. (2001). Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *Journal of ethnopharmacology*, 74(2), 113-123.
- Aketch, C.A. (1992). A preliminary survey of the conservation status of some plant species in Kenya. In: Edwards, S., Asfaw, Z. (Eds.), *The Status of Some Plant Resources in Parts of Tropical Africa. Botany 2000: East and Central Africa. NAPRECA Monograph Series No. 2. NAPRECA, Addis Ababa University, Addis Ababa, 57–65.*
- Ali, D., Hussein, S.M.S., Malik, A., Ahmed, Z. (2003). Chemical constituents of the genus *Launaea*. *Journal of Chemical Society of Pakistan*, 25, 341-347
- Apu, A. S., Muhit, M. A., Tareq, S. M., Basak, D., & Islam, M. S. (2010). Isolation and identification of compounds from the leaf extract of *Dillenia indica* Linn. *Bangladesh Pharmaceutical Journal*, 13(1), 49-53.
- Arokiyaraj, S., Sripriya, N., Bhagya, R., Radhika, B., Prameela, L., Udayaprakash, N.K. (2012). High rates of multidrug resistance in *Enterococcus faecalis* and *E. faecium* isolated from inpatients and outpatients in Taiwan. *Journal of Diagnostic Microbiology and Infectious Disease*, 75, 406–411
- Asano, S.I., Nukumizo, Y., Bando, H., Hzuka, T., Yamamoto, T., (1997). Cloning of novel enterotoxin genes from *Bacillus cereus* and *Bacillus thuringiensis*. *Applied Environmental Microbiology*, 63 (3), 1054– 1057.
- Aseer, M., Sujith, S., Seghal, K., Joseph, S., Chippu, S., 2009. Cytotoxic Potentials of Red Alga, *Laurencia brandenii* Collected from the Indian Coast. *Global Journal of Pharmacology*, 3(2), 90-94
- Ashafa, A. O. T., Orekoya, L.O., Yakubu, M.T., (2012). Toxicity profile of ethanolic extract of *Azadirachta indica* stem bark in male Wistar rats. *Asian Pacific Journal of Tropical Biomedicine*, 813-817
- Assogbadjo, A.E., Glèlè Kakaï, R., Chadare, F.J., Thomson, L., Kyndt, T., Sinsin, B., Van Damme, P., (2008). Folk classification, perception and preferences of baobab products in West Africa: Consequences for species conservation and improvement. *Journal of Economic Botany*, 62, 74–84.
- Bamford, M.K, Henderson, Z.L. (2003). A reassessment of the wooden fragment from Florisbad, South Africa. *Journal of Archaeological Science*, 30, 637–651

- Beentje, H.J. (1994). Kenya trees, shrubs and lianas. National Museums of Kenya, Nairobi, Kenya. 722
- Bennik, M. H. J., Van Overbeek, W., Smid, E. J., & Gorris, L. G. M. (1999). Biopreservation in modified atmosphere stored mungbean sprouts: the use of vegetable-associated bacteriocinogenic lactic acid bacteria to control the growth of *Listeria monocytogenes*. *Letters in applied microbiology*, 28(3), 226-232.
- Byarugaba, D.K., (2004). A view on antimicrobial resistance in developing countries and responsible risk factors. *International Journal of Antimicrobial Agents*, 24, 105–110.
- Cheng, F. W., Kai, Y., Hai, M.Z., Jie, C., Rui, F., Zhi, L. L., Shu, S.D., Yong. Y.W., Zhi, W.D., Ligang. Z. (2011). Components and Insecticidal Activity against the Maize Weevils of *Zanthoxylum schinifolium* Fruits and Leaves, *Molecules*, 16, 3077-3088.
- Chrian, M., Erasto, P., Otieno, N.J. (2011). Antimycobacterial activity and cytotoxicity effects of extracts of *Hallea rubrostipulata* and *Zanthoxylum chalybeum*. *Spatula DD* 1(3), 147-159.
- Christian, Z., (2008). Sesquiterpene lactones and their precursors as chemosystematic markers in the tribe Cichorieae of the Asteraceae. *Phytochemistry*, 69, 2270–2296.
- Christian, Z & Vipaporn, S., (2010). Flavonoids as chemosystematics markers in the tribe Cichorieae of the Asteraceae. *Journal of Biochemical Systematics and Ecology*, 38, 935–957.
- Čížek, A., Dolejská, M., Novotná, R., Haas, D., & Vyskočil, M. (2008). Survey of Shiga toxigenic *Escherichia coli* O157 and drug-resistant coliform bacteria from in-line milk filters on dairy farms in the Czech Republic. *Journal of applied microbiology*, 104(3), 852-860.
- Clinical and Laboratory Standards Institute (CLSI), (2006). Performance Standards for Antimicrobial Disk Susceptibility Tests; Approved Standard—Ninth Edition. Clinical and Laboratory Standards Institute document M2-A9 [ISBN 1-56238-586-0]. Clinical and Laboratory Standards Institute, 940 West Valley Road, Suite 1400, Wayne, Pennsylvania 19087-1898 USA.
- Clinical and Laboratory Standards Institute (CLSI), (2008). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts: Approved Standard M27-A3, NCCLS, Wayne, PA.
- Clinical and Laboratory Standards Institute (CLSI), (2009). Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically: Approved Standard, eighth ed. M07-A8. CLSI, Wayne, PA.
- Compaoré, W.R., Nikiéma, P.A., Bassolé, H.I.N., Savadogo, A., Mouecoucou, J., Hounhouigan, D.J., Traoré, S.A. (2011). Chemical Composition and Antioxidative Properties of Seeds of *Moringa oleifera* and Pulps of *Parkia biglobosa* and *Adansonia digitata* Commonly used in Food Fortification in Burkina Faso. *Current Research Journal of Biological Sciences*, 3(1), 64-72.

- Cook, D. J., Walter, S. D., Cook, R. J., Griffith, L. E., Guyatt, G. H., Leasa, D., & Brun-Buisson, C. (1998). Incidence of and risk factors for ventilator-associated pneumonia in critically ill patients. *Annals of internal medicine*, 129(6), 433-440.
- Cotton, C.M., (1996). *Ethnobotany: Principles and Applications*. Wiley, Chichester.
- Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical microbiology reviews*, 12(4), 564-582.
- Cravioto, A., Tell, A., Navarro A., (1991). Association of Escherichia coli HEp-2 adherence patterns with type and duration of diarrhea. *Lancet infectious diseases*, 337, 262-264.
- Daboor, S. M., & Haroon, A. M. (2012). *In vitro*: Antimicrobial potential and phytochemical screening of some Egyptian aquatic plants. *The Egyptian Journal of Aquatic Research*, 38(4), 233-239.
- Das, K., Tiwari, R.K.S., & Shrivastava, D.K., (2010). Techniques for Evaluation of Medicinal Plant Products as Antimicrobial Agent: Current Methods and Future Trends. *Journal of Medicinal Plants Research*, 4(2), 104-111.
- Davis, R. J. (1994). MAPKs: new JNK expands the group. *Trends in biochemical sciences*, 19(11), 470-473.
- Denis, B & Ragai, K.I., (1996). Isoprenylated flavonoids--a Survey. *Phytochemistry*, 4(5), 921-982
- DiazGranados, C.A., Denise M. Cardo D.M., McGowan Jr., J.E., (2008). Antimicrobial resistance: international control strategies, with a focus on limited-resource settings. *International Journal of Antimicrobial Agents*, 32, 1-9
- Durmaz, B., Durmaz, R., Sahin, K. (1997). Methicillin-resistance among Turkish isolates of *Staphylococcus aureus* strains from nosocomial and community infections and their resistance patterns using various antimicrobial agents. *Journal of Hospital Infection*, 37, 325-329
- Drobniewski, F.A. (1993) *Bacillus cereus* and related species. *Journal of Clinical Microbiology*, 6, 324-38.
- Edelstein, M.V., Skleenova, E.N., Shevchenko, O.V., D'souza, J.W., Tapalski, D.V., Azizov, I. S. (2013). Spread of extensively resistant VIM-2-positive ST235 *Pseudomonas aeruginosa* in Belarus, Kazakhstan, and Russia: a longitudinal epidemiological and clinical study. *Lancet Infectious Disease*, 10, 867-76.
- Essawi, T., Srour, M. (2000). Screening of some Palestinian medicinal plants for antibacterial activity, *Journal of Ethnopharmacology*, 70, 343-349.

- Fairouz, M., Amar, Z., Narimane, S., Ahmed, T., Salah, R.(2010). Flavonoid Constituents from Algerian *Launaea resedifolia* (O.K.) and Their Antimicrobial Activity. *Natural Product*, 4(1), 91-95
- Finney D.J., (1971). Probit analysis, 3rd Ed, Cambridge University press.
- Gbadamosi, I.T., Moody J. O., Lawal, A. M. (2011). Phytochemical screening and proximate analysis of eight ethnobotanicals used as antimalaria remedies in Ibadan. *Nigeria Journal of Applied Biosciences*, 44, 2967 – 2971
- Gislaine F. Moura-Costa, Samara R. Nocchi, Ligia F. Ceole, Jo~ao Carlos P. de Mello, Celso Vataru Nakamura, Benedito Prado Dias Filho, Livia G. Temponi, Tania Ueda-Nakamura, (2012). Phytochemical Screening, Antibacterial and Free Radical Scavenging Effects of *Artemisia nilagirica*, *Mimosa pudica* and *Clerodendrum siphonanthus*. An *in-vitro* study. *Asian Pacific Journal of Tropical Biomedicine*, S601-S604
- Gilbert, R. J. (1979). *Bacillus cereus* gastroenteritis. In Foodborne Infections and Intoxications, 2nd ed. (H. Riemann and F. L. Bryan). NewYork
- Graf, B.M., Martin E. (2000). The intensive Care Physician and Control of Antimicrobial Resistance. *International Journal of Antimicrobial Agents*, 16, 511–514
- Hamed, O., (2011). Studies Toward Isolation and Identification of Bioactive Substances from Medicinal Plants (Doctoral dissertation, National University, Palestine).
- Harborne, J.B., (2002). Phytochemical Methods, A Guide to Modern Techniques of Plant Analysis, 3rd Ed. Chapman & Hall. London
- He, W., Van Puyvelde, L., De Kimpe, N., Verbruggen, L., Anthonissen, K., Van der Flaas ,M., Bosselaers, J., Mathenge, S.M., Mudida,F.P.(2002) Chemical Constituents and Biological Activities of *Zanthoxylum usambarense*. *Phytotherapy Research*, 16, 66–70.
- Hood, R.D., (2009). The Importance of Toxicity Threshold for Biomonitoring, University of Albana. Retrieved from www.alb.ac.ke 22/06/2013
- Jerry, L.M. & Lingling, L.R., 1998, The Use of Biological Assays to Evaluate Botanicals. *Drug information Journal*, 32, 513-524
- Kalayou, S., Haileselassie, M., Gebre-egziabher, G., Tiku', T., Sahle, S.,Taddele, H., Ghezu, M. (2012). In-vitro antimicrobial activity screening of some ethnoveterinary medicinal plants traditionally used against mastitis, wound and gastrointestinal tract complication in Tigray Region, Ethiopia. *Asian Pacific Journal of Tropical Biomedicine*, 2(7), 512-522
- Kamuhabwa,A., Nshimo,C.,Witte,P. (2000),Cytotoxicity of some medicinal plant extracts used in Tanzanian traditional medicine, *Journal of Ethnopharmacology*, 70, 143–149
- Kanafani, Z. A., and Fowler Jr.V.G. (2006).*Staphylococcus aureus* Infections: New Challenges from an Old Pathogen. *Infeciosas y Microbiologia Clinica*, 24,182-193

- Kaper, J.B., Nataro, J.P., Mobley, H.L. (2004). Pathogenic *Escherichia coli*. *Nature Reviews Microbiology*, 2, 123–140
- Karem, Z., German-Shashoua, H., & Yarden, O., (2005). Microwave-assisted extraction of bioactive saponins from chickpea (*Cicer arietinum* L.). *Journal of the Science of Food and Agriculture*, 85, 406–412
- Kareru, P. G., Kenji, G. M., Gachanja, A. N., Keriko, J. M., Mungai, G.(2007). Traditional medicines among the Embu and Mbeere peoples of Kenya. *African. Journal of traditional, Complimentary and alternative medicine*, 4(1), 75 – 86
- Karou, D., Savadogo, A., Canini, A., Yameogo, S., Montesano, C., Simpore, J., ... Traore, A. (2006). Antibacterial activity of alkaloids from *Sida acuta*. *African Journal of Biotechnology*, 5(2), 195-200.
- Katariina, V. M. (2000). World vegetables used by the Sambia in the Usambara Mountains, NE Tanzania. *Annales Botanici Fennici*, 37, 57-67
- Kato, A., Moriyasu, M., Ichimaru, M., Nishiyama, Y., Juma, F.D., Nganga, J.N., Mathenge, S.G., Ogeto, J.Q. (1996). Isolation of alkaloidal constituents of *Zanthoxylum usambarense* and *Zanthoxylum chalybeum* using Ion-Pair HPLC. *Journal of Natural Products*, 59, 316–318.
- Keter, K.L., Mutiso, C. P. (2011). Ethnobotanical studies of medicinal plants used by Traditional Health Practitioners in the management of diabetes in Lower Eastern Province, Kenya. *Journal of Ethnopharmacology*, 1-7
- Kitonde, C. K., Fidahusein, D. S., Lukhoba, C. W., & Jumba, M. M. (2013). Antimicrobial activity and phytochemical study of *vernonia glabra* (steetz) oliv. & hiern. in Kenya. *African Journal of Traditional, Complementary and Alternative Medicines*, 10(1), 149-157.
- Kokwaro, J.O.,(2009). Medicinal plants of East Africa, Nairobi University Press. 3rd Edition. Nairobi
- Kone, W. M., K. K. Atindehou, C. Terreaux, K. Hostettmann, D. Traore, and M. Dosso. 2004. Traditional medicine in North Cote-d'Ivoire: screening of 50 medicinal plants for antibacterial activity. *Journal of Ethnopharmacology*, 93, 43-49.
- Koua, F.H.M., Babiker, H.A., Halfawi, A., Ibrahim, R.O., Abbas, F.M., Elgaali, E.I., ... Khlafallah, M.M. (2011). Phytochemical and Biological Study of *Striga hermonthica* (Del.) Benth Callus and Intact Plant. *Research In Pharmaceutical Biotechnology*, 3(7), 85-92.
- Krishanaraju, A. V., Rao, T. V., Sundararaju, D., Vanisree, M., Tsay, H. S., & Subbaraju, G. V. (2005). Assessment of Bioactivity of Indian Medicinal Plants Using Brine Shrimp (*Artemia Salina*) Lethality Assay+ .
- Lyimo, M., Temu, R.P.C., Mugula, J.K. (2003). Identification and Nutrient Composition of Indigenous Vegetables of Tanzania. *Plant Foods for Human Nutrition*, 58, 85–92

- Manilal, A., Sujith, S., Kiran, G. S., Selvin, J., & Shakir, C. (2009). Cytotoxic potentials of red alga, *Laurencia brandenii* collected from the Indian coast. *Global Journal of Pharmacology*, 3(2), 90-94.
- Marchese A., Gualco L., Debbia E.A., Bandettini R., Pescetto L., Cavallero A., Ossi M.C., Schito A.M. (2007). Antifungal resistance in *Candida spp.* isolated in Italy between 2002 and 2005 from children and adults. *International Journal of Antimicrobial Agents*, 29, 179–184
- Mariita, R.M., Ogol, C.K.P., Ogege, N.O., & Okemo, P.O. (2010). Antitubercular and Phytochemical Investigation of Methanol Extracts of Medicinal Plants used by the Samburu Community in Kenya. *Tropical Journal of Pharmaceutical Research*, 9(4), 379-385.
- Marimuthu, J., Mithraja, M.J., Mony Mahesh, M., Paul, Z. M., Jeeva, S., (2011). Phytochemical studies on *Azolla pinnata* R. Br., *Marsilea minuta* L. and *Salvinia molesta* Mitch. *Asian Pacific Journal of Tropical Biomedicine* S26-S29
- Maryam, Z., Aqil, F., Khan, M.S.A., & Ahmad, I. (2010). Ethnomedicinal plants derived antibacterial and their prospects. *Ethnomedicine. A source of Complementary Therapeutics*, 150-177.
- Masakazu, N., Firth, N.A., Cannon, D.R. (2010). Antifungal drug resistance of oral fungi. *Odontology*, 98(1), 15-25.
- Matu, E.N., VanStaden, J. (2003). Antibacterial and anti-inflammatory activities of some plants used for medicinal purposes in Kenya. *Journal of Ethnopharmacology* 87, 35–41
- Mazid, M.A., Datta, B.K., Nahar, L., Sarker, S.D. (2008). Assessment of anti-bacterial activity and brine shrimp toxicity of two *Polygonum* species. *Ars Pharmaceutica*, 49(2), 127–134.
- Mazid, M. , Khan, T.A., Mohamad, F., (2011). Role of Secondary Metabolites in Defence Mechanism in Plants. *Journal of Biology and Medicine*, 3(2), 232-249
- McCarthy, J.G., (1990). *Plastic surgery*, Philadelphia: Saunders, 0-7216- 1514-7, p. 807–808.
- McEwen, S. A., & Fedorka-Cray, P. J. (2002). Antimicrobial use and resistance in animals. *Clinical Infectious Diseases*, 34(Supplement 3), S93-S106.
- McGaw, L. J., Rabe, T., Sparg, S. G., Jäger, A. K., Eloff, J. N., & Van Staden, J. (2001). An investigation on the biological activity of *Combretum* species. *Journal of ethnopharmacology*, 75(1), 45-50.
- Meyer, B.N., Ferrigni, N.R., Putnam, J.E., Jacobsen, J.B., Nicholsand, D.E., Mclaughlin, J.L. (1982). Brine shrimp; a convenient general bioassay for active plant constituents, *Planta Medica*, 45, 31–34.

- Miaron, O.J., Kassim, O., & Ekaya, N. (2004). Indigenous Knowledge: The basis of Maasai Ethnoveterinary Diagnostic Skills. *Journal of Human Ecology*, 16, 43-48.
- Mubarak, S., Robert, A.A., Baskaradoss, J.K., Al-Zoman, K., Al Sohail, A., Alsuwyed, A. (2013). The prevalence of oral *Candida* infections in periodontitis patients with type 2 diabetes mellitus. *Journal of Infection and Public Health*, 6(4), 296-301
- Munguti, K. (1997). Indigenous Knowledge in the Management of Malaria and Visceral Leishmaniasis among the Tugen of Kenya. *Indigenous Knowledge Development Monitor*, 5, 10-12
- Musila, M.F, Dossaji, S.F., Nguta, J.M., Lukhoba, C.W., Munyao, J.M. (2013). *In vivo* antimalarial activity, toxicity and phytochemical screening of selected antimalarial plants. *Journal of Ethnopharmacology*, 146, 557-561
- Nagata , J.M., Alan, R. J., John, .K. Charles,R.S.,Elizabeth,A.B.,Craig,R.C. (2011). Medical pluralism on Mfangano Island: Use of medicinal plants among persons living with HIV/AIDS in Suba District, Kenya. *Journal of Ethnopharmacology*, 135, 501–509
- Nester, E.W., Anderson, D.G., Roberts, C.E., Pearsall, N.N., & Nester, M.T. (2004). *Microbiology: A Human Perspective*. 4th ed, McGraw-Hill. New York..
- Nguta, J.M., Mbaria, J.M., Gakuya, D.W., Gathumbi, P.K., Kiama, S.G. (2010). Antimalarial herbal remedies of Msambweni, Kenya. *Journal of Ethnopharmacology*, 128, 424–432
- Nguta,J.M., Mbaria,J.M., Gakuya,D.W.,Gathumbi,P.K.,Kabasa,J.D.,Kiama,S.G. (2011). Biological Screening of Kenya Medicial Plants using *Artemia salina* L. (Artemiidae), *Pharmacology online*, 2, 458-478
- Nizet , V., Bradley, J.S., (2011). Chapter 14: *STAPHYLOCOCCAL* INFECTIONS, Bacterial infections, section 1,489-515 DOI: 10.1016/B978-1-4160-6400-8.00014-6
- Njoroge, G. N., & Bussmann, R. W. (2007). Ethnotherapeutic management of skin diseases among the Kikuyus of Central Kenya. *Journal of ethnopharmacology*, 111(2), 303-307.
- Olila, D., Olwa-Odyek., Opuda-Asibo.J., (2002). Screening of extracts of *Zanthoxylumchalybeum* and *Warburgia ugandensis* for activity against measles virus (Swartz and Edmonston strains) in vitro. *African Health Sciences*, 2, 2-10
- Paphitou, N.I.,(2013). Antimicrobial resistance: action to combat the rising microbial challenges. *International Journal of Antimicrobial Agents*, article in press
- Pesti, M., Virág, E., Belagyi, J., Gazdag, Z., Vágvölgyi, C. (2012). Direct in vivo interaction of the antibiotic primycin with the plasma membrane of *Candida albicans*: An EPR study. *Biochimica et Biophysica Acta*, 1818, 42-48
- Pirttijärvi T.S.M., Andersson M.A., Scoging A.C., Salkinoja-Salonen M.S. (1999). Evaluation of methods for recognizing strains of the *Bacillus cereus* group with food poisoning

- potential among industrial and environmental contaminants, *System. Applied Microbiology*, 22, 133–144.
- Raghunathaiyar, S., (1996). *Indian Medicinal Plants* 3, 104–105.
- Ramadan, A., Harraz, F.M., EL-Mougy, S.A.,(1994). Anti-inflammatory, analgesic and antipyretic effects of the fruit pulp of *Adansonia digitata*, *Fitoterapia*, 65(5), 418-422
- Rasoanaivo, P., Ratsimamanga-Urveg, S., (1993). Biological Evaluation of Plants with Reference to the Malagasy Flora. Monograph of the IFS-NAPRECA Workshop on Bioassays, Antananarivo, Madagascar, 72–79.
- Rathee, D., Rathee ,D., Rathee, P., Rathee, S. (2012). Phytochemical screening and antimicrobial activity of *Picrorrhiza kurroa*, an Indian traditional plant used to treat chronic diarrhea. *Arabian Journal of Chemistry*, in press
- Ross, J. A., & Kasum, C. M. (2002). Dietary flavonoids: bioavailability, metabolic effects, and safety. *Annual review of Nutrition*, 22(1), 19-34.
- Rukunga, G.M., Gathirwa, J.W., Omar, S.A., Muregi, F.W., Muthaura, C.N., Kirira, P.G., Mungai, G.M., Kofi-Tsekpo, W.M. (2009). Anti-plasmodial activity of the extracts of some Kenyan medicinal plants. *Journal of Ethnopharmacology*, 121, 282–285
- Sam, T.W., (1993). Toxicity testing using the brine shrimp: *Artemia salina*. In: Colegate S.M., Molyneux, R.J., editors. *Bioactive natural products detection, isolation, and structural determination*. Boca Raton: CRC press, 442–456
- Sandiga, I., Chacha, C.N., & Kanunah, M.P. (1995). *Traditional Medicine in Africa: The existence and Use of Traditional Medicine in Kenya* [e-book]. Nairobi: East African Educational Publishers Ltd. Available online: <http://books.google.com/books> [accessed 20th January 2014].
- Schultz, J. C., Hunter, M. D., & Appel, H. M. (1992). Antimicrobial activity of polyphenols mediates plant-herbivore interactions. *In Plant polyphenols*, 621-637
- Shahat, A.A.,2006, Procyanidins from *Adansonia digitata*. *Journal of Pharmaceutical Biology*, 44(6), 445–450
- Shimeld, L.A., and Rodgers A.T., (1999). *Essentials of Diagnostic Microbiology*. New York: Delmar publishers
- Sidibe, M., William, J.T. (2002). *BAOBAB: Adansonia digitata L.*, International Center for Underutilized Crops, RPM Reprographics Chichester, England.
- Singh, H. B., & Handique, A. K. (1997). Antifungal Activity of the Essential Oil of *Hyptis suaveolens* and Its Efficacy in Biocontrol Measures. *Journal of Essential Oil Research*, 9(6), 683-687.

- Shankar, T. B., Shantha, N. V., Ramesh, H. P., Murthy, I. A., & Murthy, V. S. (1980). Toxicity studies on turmeric (*Curcuma longa*): acute toxicity studies in rats, guineapigs and monkeys. *Indian journal of experimental biology*, 18(1), 73-75.
- Shankar, S. R., Rangarajan, R., Sarada, D. V. L., & Kumar, C. S. (2010). Evaluation of Antibacterial Activity and Phytochemical Screening of *Wrightia tinctoria* L. *Pharmacognosy Journal*, 2(14), 19-22.
- Sher, A. (2004). Antimicrobial activity of natural products from medicinal plants. *Gomal Journal of Medical Sciences*, 7(1), 1-5
- Sleet, R.B., Brendel, K. (1983). Improved methods for harvesting and counting synchronous populations of *Artemia nauplii* for use in developmental toxicology. *Ecotoxicology and Environmental Safety*, 7, 435–446.
- Sukumaran, S., Kiruba, S., Mahesh, M., Nisha, S.R., Miller, P. Z., Ben, C.P., Jeeva, S. (2011). Phytochemical constituents and antibacterial efficacy of the flowers of *Peltophorum pterocarpum*(DC.) Baker ex Heyne. *Asian Pacific Journal of Tropical Medicine*, 735-738
- Takate, S.B., Pokharkar, R.D., Chopade, V.V., Gite, V.N., (2010). Hepato-protective activity of the ethyl acetate extract of *Launaea intybacea* (Jacq) Beauv in paracetamol induced hepato-toxicity in albino rats, *International Journal of Pharmaceutical Science*, 1(2), 72-74
- Taniguchi, M., Chapya, A., Kubo, I., Nakanishi, K., (1978). Screening of East African plants for antimicrobial activity I. *Chemical and Pharmaceutical Bulletin*, 26, 2910–2913
- Tollefson, L., and Miller M. A., (2000). Antibiotic use in food animals: controlling the human health impact. *International journal of food microbiology*, 83, 245-254.
- Touitou, I., Lesage, S., McDermott, M., Cuisset, L., Hoffman, H., Dode, C., & de Menthiere, C. S. (2004). Infevers: an evolving mutation database for auto-inflammatory syndromes. *Human mutation*, 24(3), 194-198.
- United Nations Environment Program- UNEP. (2010). Global Strategy for Plant Conservation. Biodiversity is life 2010 International Year of Biodiversity. Online: www.cbd.int/gspc [accessed 20th January, 2014]
- Vanhaecke, P., Persoone, G., Claus, C., Sorgeloos, P. (1981). Proposal for a short-term toxicity test with *Artemia nauplii*. *Ecotoxicology and Environmental Safety*, 5, 382–387
- VanStaden, J, Mulaudzi, R.B., Ndhlala, A.R., Kulkarni, M.G., Finnie, J.F. (2012). Anti-inflammatory and mutagenic evaluation of medicinal plants used by Venda people against venereal and related diseases. *Journal of Ethnopharmacology*, 146, 173-179.
- VanStaden J., Mulaudzi, R.B., Ndhlala, A.R., Kulkarni, M.G., Finnie, J.F. (2011). Antimicrobial properties and phenolic contents of medicinal plants used by the Venda people for conditions related to venereal diseases. *Journal of Ethnopharmacology*, 137, 330-337

- Vazquez, J.A. (1999). Options for the management of mucosal candidiasis in patients with AIDS and HIV infection. *Pharmacotherapy*, 19, 76–87
- Vital, P.G., and Rivera, W.L., (2011). Antimicrobial activity, cytotoxicity, and phytochemical screening of *Voacanga globosa* (Blanco) Merr. leaf extract (Apocynaceae). *Asian Pacific Journal of Tropical Medicine*, 824-828
- Wagner, H., (1997). Guidelines for quality control of phytomedicines. In: World Congress on Medicinal and Aromatic Plants for Human Welfare, vol. 2, Abstracts. ICMIPA/ISHS/SAIPOA, Mendoza, p. L-4
- Waksmundzka, H.M. , Sherma, J., Kowalska, T. (2008). Thin Layer Chromatography in Phytochemistry, Chromatography Science Series 99.CRC Press
- Wang, Y.F., Ni, Z. Y., Dong, M., Cong, B., Shi, W. Q., Gu, Y. C., Kiyota, H. (2010). Secondary Metabolites of Plants from the Genus *Saussurea*; Chemistry and Activity. *Chemistry and Biodiversity*, 7, 2623-2659
- Wanyoike, G.N., Chabra, S.C., Lang’at, C.C., Omar, S.A., (2004). Brine shrimp toxicity and antiplasmodial activity of five Kenyan medicinal plants. *Journal of Ethnopharmacology*, 90, 129–133
- Weber, D. J., Saviteer, S. M., Rutala, W. A., and Thomann, C. A., (1988). In vitro susceptibility of *Bacillus* spp. to selected antimicrobial agents. *Antimicrobial Agents and Chemotherapy*, 32, 642- 645.
- Williams RP.(1986). *Bacillus anthracis* and other- spore-forming bacilli. In: Braude Al, Davis (CE, Fierer J, eds. *Infectious Diseases and Medical Microbiology*. 2nd ed. Philadelphia: W.B. Saunders; 1514-1518.
- Whiting, M. F. (2001). Phylogeny of the holometabolous insect orders: molecular evidence. *Zoologica Scripta*, 31(1), 3-15.
- World Health Organization (WHO). 2001. Legal status of traditional medicine and complementary/alternative medicine: a worldwide review. Available from: http://libdoc.who.int/hq/2001/WHO_EDM_TRM_2001.2.pdf
- World Health Organization, Food Agricultural Organization of the United Nations and World Organization for Animal Health- W.H.O/FAO/OIE. (2003). Expert Workshop on Non-human Antimicrobial Usage and Antimicrobial Resistance: *Scientific Assessment*, 1-5
- Yusuf, S., Mehta, S. R., Peters, R. J., Bertrand, M. E., Lewis, B. S., Natarajan, M. K., ... & Fox, K. A. (2001). Effects of pretreatment with clopidogrel and aspirin followed by long-term therapy in patients undergoing percutaneous coronary intervention: the PCI-CURE study. *The Lancet*, 358(9281), 527-533.

APPENDICES

Appendix 1: Growth inhibition of *B. cereus* by aqueous extracts

Growth inhibition of *B. cereus* at 200mg/ml
Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-7.16667*	.36004	.000	-8.2112	-6.1221
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-5.33333*	.36004	.000	-6.3779	-4.2888
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-15.83333*	.36004	.000	-16.8779	-14.7888
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-15.83333*	.36004	.000	-16.8779	-14.7888
DMSO Water	& Gentamicin(40ug/ml)	-15.83333*	.36004	.000	-16.8779	-14.7888

*. The mean difference is significant at the 0.05 level.

Growth inhibition of *B. cereus* at 100mg/ml
Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-8.33333*	.31914	.000	-9.2592	-7.4074
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-8.16667*	.31914	.000	-9.0926	-7.2408
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-15.83333*	.31914	.000	-16.7592	-14.9074
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-15.83333*	.31914	.000	-16.7592	-14.9074
DMSO & Water	Gentamicin(40ug/ml)	-15.83333*	.31914	.000	-16.7592	-14.9074

*. The mean difference is significant at the 0.05 level.

Appendix 2: Growth inhibition of *B. cereus* by organic extracts

Growth inhibition of *B. cereus* at 200 mg/ml

Dunnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-1.00000	.43033	.000	-9.4001	-4.9333
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-2.16667	.76980	.058	-4.4001	.0667
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-4.00000*	.76980	.001	-6.2334	-1.7666
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-15.83333*	.76980	.000	-18.0667	-13.5999
DMSO Water	& Gentamicin(40ug/ml)	-15.83333*	.76980	.000	-18.0667	-13.5999

*. The mean difference is significant at the 0.05 level.

Growth inhibition of *B. cereus* at 100 mg/ml

Dunnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-8.66667*	.67358	.000	-10.6209	-6.7125
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-3.66667*	.67358	.001	-5.6209	-1.7125
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-15.83333*	.67358	.000	-17.7875	-13.8791
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-6.16667*	.67358	.000	-8.1209	-4.2125
DMSO Water	& Gentamicin(40ug/ml)	-15.83333*	.67358	.000	-17.7875	-13.8791

*. The mean difference is significant at the 0.05 level.

Appendix 3: Growth inhibition MRSA by aqueous extracts

Growth inhibition of MRSA at 200 mg/ml

Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-7.16667*	0.76980	.000	-2.2485	.2485
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-15.33333*	.43033	.000	-16.5818	-14.0848
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-15.33333*	.43033	.000	-16.5818	-14.0848
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-15.33333*	.43033	.000	-16.5818	-14.0848
DMSO Water	& Gentamicin(40ug/ml)	-15.33333*	.43033	.000	-16.5818	-14.0848

*. The mean difference is significant at the 0.05 level.

Growth inhibition of MRSA at 100 mg/ml

Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-3.33333*	.50918	.000	-4.8106	-1.8561
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-15.33333*	.50918	.000	-16.8106	-13.8561
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-15.33333*	.50918	.000	-16.8106	-13.8561
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-15.33333*	.50918	.000	-16.8106	-13.8561
DMSO Water	& Gentamicin(40ug/ml)	-15.33333*	.50918	.000	-16.8106	-13.8561

*. The mean difference is significant at the 0.05 level.

Appendix 4: Growth inhibition of MRSA by the organic extracts

Growth inhibition of MRSA at 200 mg/ml

Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(0.04/ml)	-5.66667*	.66667	.055	-7.6008	-3.7325
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-5.66667*	.66667	.000	-7.6008	-3.7325
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-7.00000*	.66667	.000	-8.9342	-5.0658
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-4.66667*	.66667	.000	-6.6008	-2.7325
DMSO Water	& Gentamicin(40ug/ml)	-15.00000*	.66667	.000	-16.9342	-13.0658

*. The mean difference is significant at the 0.05 level.

Growth inhibition of MRSA at 100 mg/ml

Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-6.83333*	.44096	.000	-8.1127	-5.5540
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-5.66667*	.44096	.000	-6.9460	-4.3873
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-7.50000*	.44096	.000	-8.7793	-6.2207
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-5.83333*	.44096	.000	-7.1127	-4.5540
DMSO Water	& Gentamicin(40ug/ml)	-15.00000*	.44096	.000	-16.2793	-13.7207

*. The mean difference is significant at the 0.05 level.

Appendix 5: Growth inhibition of *P. aeruginosa* by aqueous extracts

Growth inhibition of *P. aeruginosa* at 200 mg/ml
Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-9.33333*	.50918	.000	-10.8106	-7.8561
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-18.50000*	.50918	.000	-19.9772	-17.0228
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-18.50000*	.50918	.000	-19.9772	-17.0228
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-18.50000*	.50918	.000	-19.9772	-17.0228
DMSO Water	& Gentamicin(40ug/ml)	-18.50000*	.50918	.000	-19.9772	-17.0228

*. The mean difference is significant at the 0.05 level.

Growth inhibition of *P. aeruginosa* at 100 mg/ml
Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-10.50000*	.55277	.000	-12.1037	-8.8963
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-18.50000*	.55277	.000	-20.1037	-16.8963
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-18.50000*	.55277	.000	-20.1037	-16.8963
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-18.50000*	.55277	.000	-20.1037	-16.8963
DMSO Water	& Gentamicin(40ug/ml)	-18.50000*	.55277	.000	-20.1037	-16.8963

*. The mean difference is significant at the 0.05 level.

Appendix 6: Growth inhibition of *P. aeruginosa* by organic extracts

Growth inhibition of *P. aeruginosa* at 200 mg/ml

Dunnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-9.33333*	.72008	.000	-11.4225	-7.2442
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-8.00000*	.72008	.000	-10.0891	-5.9109
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-8.00000*	.72008	.000	-10.0891	-5.9109
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-7.83333*	.72008	.000	-9.9225	-5.7442
DMSO & Water	Gentamicin(40ug/ml)	-18.50000*	.72008	.000	-20.5891	-16.4109

*. The mean difference is significant at the 0.05 level.

Growth inhibition of *P. aeruginosa* at 100 mg/ml

Dunnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Gentamicin(40ug/ml)	-9.50000*	.60093	.000	-11.2434	-7.7566
<i>Z. chalybeum</i>	Gentamicin(40ug/ml)	-9.33333*	.60093	.000	-11.0768	-7.5899
<i>L. cornuta</i>	Gentamicin(40ug/ml)	-9.33333*	.60093	.000	-11.0768	-7.5899
<i>G. trichocarpa</i>	Gentamicin(40ug/ml)	-8.83333*	.60093	.000	-10.5768	-7.0899
DMSO & Water	Gentamicin(40ug/ml)	-18.50000*	.60093	.000	-20.2434	-16.7566

*. The mean difference is significant at the 0.05 level.

Appendix 7: Growth inhibition of *C. albicans* by organic extracts

Growth inhibition of *C. albicans* at 200 mg/ml

Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Amphotericin (30ug/ml)	-6.66667*	.36004	.000	-7.7112	-5.6221
<i>Z. chalybeum</i>	Amphotericin (30ug/ml)	-7.50000*	.36004	.000	-8.5446	-6.4554
<i>L. cornuta</i>	Amphotericin (30ug/ml)	-15.66667*	.36004	.000	-16.7112	-14.6221
<i>G. trichocarpa</i>	Amphotericin (30ug/ml)	-7.16667*	.36004	.000	-8.2112	-6.1221
DMSO & Water	Amphotericin (30ug/ml)	-15.66667*	.36004	.000	-16.7112	-14.6221

*. The mean difference is significant at the 0.05 level.

Growth inhibition of *C. albicans* at 100 mg/ml

Dunnnett t (2-sided)

(I) Treatments	(J) Treatments	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
					Lower Bound	Upper Bound
<i>A. digitata</i>	Amphotericin (30ug/ml)	-7.83333*	.28868	.000	-8.6709	-6.9958
<i>Z. chalybeum</i>	Amphotericin (30ug/ml)	-8.16667*	.28868	.000	-9.0042	-7.3291
<i>L. cornuta</i>	Amphotericin (30ug/ml)	-15.66667*	.28868	.000	-16.5042	-14.8291
<i>G. trichocarpa</i>	Amphotericin (30ug/ml)	-7.83333*	.28868	.000	-8.6709	-6.9958
DMSO & Water	Amphotericin (30ug/ml)	-15.66667*	.28868	.000	-16.5042	-14.8291

*. The mean difference is significant at the 0.05 level.

Appendix 8: Brine shrimp lethality raw data

PLANT NAME	CONCENTRATION ($\mu\text{g/ml}$)											
	1000 ($\mu\text{g/ml}$)				100 ($\mu\text{g/ml}$)				10 ($\mu\text{g/ml}$)			
	1 st trial	2 nd trial	3 rd trial	Average mortality	1 st trial	2 nd trial	3 rd trial	Average mortality	1 st trial	2 nd trial	3 rd trial	Average mortality
<i>A. digitata</i> (aq)	2	0	2	1.33	0	0	1	0.33	0	0	0	0
<i>A. digitata</i> (org)	2	2	4	2.67	1	2	0	1	0	0	0	0
<i>Z. chalybeum</i> (aq)	7	8	9	8	2	3	2	2.33	0	0	0	0
<i>Z. chalybeum</i> (org)	10	10	10	10	5	6	6	5.67	4	3	2	3
<i>L. cornuta</i> (aq)	7	6	6	6.33	2	1	2	1.67	0	0	1	0.33
<i>L. cornuta</i> (org)	8	7	10	8.33	2	4	6	4	2	0	2	1.33
<i>G. trichocarpa</i> (aq)	1	0	0	0.33	0	0	0	0	0	0	0	0
<i>G. trichocarpa</i> (org)	1	3	3	2.33	0	0	0	0	0	0	0	0