

## **Salivary human immunodeficiency virus (HIV)-1-specific immunoglobulin A in HIV-1-exposed infants in Kenya**

Farquhar, C; VanCott, T; Bosire, R; Bermudez, C; Mbori-Ngacha, DA; Lohman-Payne, B; Nduati, RW; Otieno, P; John-Stewart, G

<http://hinari-gw.who.int/whalecomwww.ncbi.nlm.nih.gov/whalecom0/pubmed/18505437>

<http://erepository.uonbi.ac.ke:8080/xmlui/handle/123456789/32168>

Date: 2008-07

### **Abstract:**

Humoral immunity, and specifically immunoglobulin A (IgA) that is directed against human immunodeficiency virus (HIV)-1, may contribute to protection against HIV-1 acquisition at mucosal surfaces. HIV-1-specific IgA has been detected in genital tract secretions of HIV-1-uninfected commercial sex workers with HIV-1 exposure, and may be produced in parotid saliva by infants exposed orally to HIV-1 during delivery and breastfeeding. To explore this hypothesis, we collected saliva from 145 infants aged  $<$  or  $=$  6 months enrolled in a perinatal HIV-1 transmission study in Nairobi and from 55 control infants without HIV-1 exposure who were born to HIV-1-seronegative mothers. Among the 145 infants, 115 (79%) remained uninfected during the 12-month study period and 30 (21%) became HIV-1-infected during follow-up. Nine (8%) of the 115 HIV-1-exposed, uninfected infants had detectable levels of HIV-1 gp160-specific IgA compared with four (13%) of 30 infected infants and none of 55 control infants ( $P = 0.47$  and  $P = 0.03$  respectively). Among the nine HIV-1-exposed, uninfected infants with positive assays, median age was 1 month and none acquired HIV-1 during follow-up. We conclude that HIV-1-specific salivary IgA responses may be generated by very young infants exposed perinatally to maternal HIV-1. Mucosal responses would be an appropriate target for paediatric vaccines against breast milk HIV-1 transmission.