

# **Observer variability of an angiographic grading scale used for the assessment of intracranial aneurysms treated with flow-diverting stents.**

## **Abstract**

### **BACKGROUND AND PURPOSE:**

Novel angiographic grading scales for the assessment of intracranial aneurysms treated with flow-diverting stents have been recently developed because previous angiographic grading scales cannot be applied to these aneurysms. The purpose of this study was to evaluate the inter- and intraobserver variability of the novel O'Kelly Marotta grading scale, which was developed specifically for the angiographic assessment of aneurysms treated with flow-diverting stents.

### **MATERIALS AND METHODS:**

Multiple raters ( $n = 31$ ) from the disciplines of neuroradiology and neurosurgery were presented with pre- and posttreatment angiographic images of 14 aneurysms treated with intraluminal flow diverters. Raters were asked to classify pre- and posttreatment angiograms by using the OKM grading scale. Statistical analyses were subsequently performed with calculation of a generalized multirater  $\kappa$  statistic for assessment of inter- and intraobserver variability and by performing a Wilcoxon signed rank sum test for assessment of group differences.

### **RESULTS:**

Variability analysis of the OKM grading scale yielded substantial ( $\kappa = 0.74$ ) and almost perfect ( $\kappa = 0.99$ ) inter- and intraobserver agreement, respectively, with no statistically significant differences between raters with a background of neuroradiology versus neurosurgery or attending physician versus trainee.

### **CONCLUSIONS:**

The OKM grading scale for the assessment of intracranial aneurysms treated with flow-diverting stents is a reliable grading scale that can be used equally well by users of varying backgrounds and levels of training. Comparison with interobserver variability of pre-existing angiographic grading scales shows equal or better performance.