



Research Article

Phytochemical screening and evaluation of antioxidant activity of methanolic extract of Kenyan *Hydnora abyssinica* A. Braun (Hydnoraceae)

Onyancha J.M^{*1}, Cherongis C.N.^{1,2}, Nzivo J.M¹, Muriithi G.I.³, Njuguna D.G⁴, Mwalukumbi J.M⁵

¹Mount Kenya University, School of Pharmacy, Department of Pharmacognosy, P.O Box 342-01000, Thika, Kenya

²East Africa Community Secretariat, P.O. Box 1096, Arusha, Tanzania

³Mount Kenya University, School of Pharmacy Department of Pharmaceutics, P.O Box 342-01000, Thika, Kenya

⁴Mount Kenya University, School of Pharmacy, Department of Pharmaceutical Chemistry, P.O Box 342-01000, Thika, Kenya

⁵University of Nairobi, School of Pharmacy, Department of Pharmacology and Pharmacognosy, P.O Box 30197-01000, Nairobi, Kenya

Abstract

Hydnora abyssinica is a leafless medicinal herb characterized by rhizome and flower only. The decoction of the rhizome has been used in Kenya to manage infections, cancer and removal of retained placenta. Plant secondary metabolites that exhibit antioxidant activity are well documented and include mainly the phenolics. The objective of this study was to carry out phytochemical screening and investigate the antioxidant activity of the methanol extract of *H. abyssinica* rhizome. The extract was prepared by maceration. Phytochemical screening was done by standard methods and *in vitro* antioxidant activity was evaluated using 1,1-diphenyl-2-picrylhydrazyl (DPPH) scavenging assay. Phytochemical studies revealed presence of alkaloids and glycosides which are reported for the first time being characteristic of this plant while tannins, phenols, steroids, flavonoids, terpenoids and fatty acids confirms reports of other similar studies on the Sudanese *Hydnora abyssinica*. Saponins, volatile oils and coumarins were not present in both the rhizome and flowers. DPPH free radical scavenging activity of the rhizome methanolic extract showed potential antioxidant activity with IC₅₀ values of 26.7 µg/ml compared with that of 29.3 µg/ml for L-ascorbic acid standard. Activity of the methanolic extract may be associated with the presence of a number of varied classes of phytochemicals. This study perhaps pave way for further studies to provide data that would validate its medicinal uses and focus on bioassay guided fractionation and isolation of active compounds from its extracts.

Key words: Radical scavenging activity, DPPH, Maceration, Phytochemical screening

***Corresponding Author:** Onyancha J.M, Mount Kenya University, School of Pharmacy, Department of Pharmacognosy, P.O Box 342-01000, Thika, Kenya. Email: jamionusus@yahoo.com

1. Introduction

Hydnora abyssinica A. (Hydnoraceae) is a leafless medicinal herb characterized by rhizome and flower only. The plant is indigenous to Africa with currently four other species of the genus *Hydnora* that have been recognized and these are *H. Africana* T., *H. Abyssinica* A., *H. esculanta* J., *H. triceps* D. and *H. Sinandevu* B. [10]. *H. abyssinica* A. synonymously referred to as *H. johannis* B.N. and *H. solmsiana* D. Sudanese traditional medical practitioners use it for curing severe bacterial infections [10,15] while in East Africa it is used as anti-diarrheal medicine, traditional medical practitioners in Kenya use the root decoction as a cure for throat complaints, as an astringent in dysentery, for treatment of stomach ache and for removing retained placenta during child birth [7]. Other reports indicate that it is used for the treatment of diarrhea, amoebic dysentery, typhoid, anthrax, East Coast Fever and cancer [9, 11] and as human and animal food [8, 10].

Natural antioxidants are secondary metabolites obtained from animal, plant, mineral or microorganism sources and prevent oxidation or formation of reactive oxygen species, such as superoxide (O_2^-), Hydroxyl (OH) and peroxy (OOH, ROO) radicals, which commonly cause oxidative stress. The reactive oxygen species play an important role related to the degenerative or pathological process of various diseases such as aging, cancer, coronary heart disease, Alzheimer's disease, neuro degenerative disorders, atherosclerosis, cataracts and inflammation [2]. This study evaluated the antioxidant and phytochemistry of *H. abyssinica* with a view of justifying its utilization in management of degenerative diseases.

2. Materials and Methods

The rhizome and flowers of *H. abyssinica* were collected in November, 2013 from Kiangombe forest in Embu county, Kenya. The plant was identified at the site of collection by local traditional medicine practitioners who also provided detailed information on folklore uses. The voucher specimen of the plant was authenticated at the Herbarium section of National Museums of Kenya where it was deposited and its duplicate was deposited at Mount Kenya University Herbarium in the School of Pharmacy with a specimen voucher number HAO-1-2013. The materials were air-dried, ground. Sequential maceration was done successfully using petroleum ether, dichloromethane, dichloromethane:methanol mixture (1:1) and methanol. The extract was filtered, reduced *in vacuo* and completely oven-dried at temperatures set at 40°C. The dried samples were stored in a freezer at - 20°C until further evaluation for antioxidant activity.

Phytochemical analysis

The powders of roots and leaves were tested for the presence of bioactive compounds using standard methods.[3, 5].

Quantitative DPPH free radical scavenging activity

A stable radical of 1,1-diphenyl -2-picrylhydrazyl (3.94 mg) was dissolved in methanol (100 ml) to give 100 μ m solutions. To 3 ml of the methanolic solutions of DPPH was added 0.5 ml of each of the methanolic extract with concentrations of 0-640 μ g/ml. The set up was kept incubated for 30 minutes in the dark and at room temperature, absorbance

(A) was measured in triplicates for each sample at 517 nm using a SHIMADZU Multispect-1501 spectrophotometer. The percentage of the radical scavenging activity (RSA) was calculated by the following equation:

$RSA\% = [A_{\text{control}} - A_{\text{sample}}] / A_{\text{control}} \times 100$ and the obtained data was analyzed using GraphPad Prism Version 5 software.

DPPH Solution (3 ml) plus methanol (0.5 ml) was used as a negative control

Methanol (3.5 ml) was used as a blank while L-Ascorbic acid at concentrations equivalent to that of the test samples (0-640 µg/ml) was used as positive control as well as a standard reference [12,17].

3. Results and Discussion

Phytochemical analysis

The results of phytochemical tests, as shown in Table 1 indicated that the rhizome and flowers of *H. abyssinica* contain alkaloids, glycosides, tannins, phenols, steroids, flavonoids, terpenoids and fatty acids while saponins, volatile oils and coumarins were not present in both rhizome and flowers. The phytochemical screening results of this Kenyan *Hydnora abyssinica* are somehow consistent with the results of Sudanese *Hydnora abyssinica* for the presence of phenols, flavonoids and tannins. In contrast with the previous report on the absence of glycosides, in this study glycosides and alkaloids were reported for the first time [15].

The presence of quite a number of classes of secondary metabolites indicated above may be predictive of the ethnomedicinal claims on this medicinal plant. These phyto- constituents may be responsible for many pharmacological actions of the plant. The alkaloids have been associated with

medicinal uses for centuries and one of their common properties is cytotoxicity but also analgesic, antispasmodic and antibacterial activities have been reported [13]. The glycosides are known to lower blood pressure and also especially cardiac glycosides have been used for over two centuries as stimulants in case of cardiac failure [13]. The anthraquinone glycosides have also been used as laxatives [13]. Steroids have antibacterial properties and they are also very important compounds especially due to their relationship with compounds such as sex hormones [13].

The phenolic compounds in this case flavonoids and tannins, possess biological properties such as anti-apoptosis, anti-aging, anti-carcinogen, anti-inflammation, anti-arteriosclerosis, cardiovascular protection and improvement of endothelial function, as well as inhibition of angiogenesis and cell proliferation activities [19]. This study of evaluation of antioxidant activity and phytochemical screening would perhaps lay a basis for further investigations based on the traditional uses of the plant in management of cancer and infectious diseases [13, 19].

Quantitative assay (*In vitro* DPPH free radical scavenging assay)

The 1, 1-diphenyl -2-picrylhydrazyl (DPPH) radical scavenging activity of *H. abyssinica* rhizome methanolic (HA-MeOH) extract compared with standard antioxidant, L-ascorbic acid (ASA-Std) showed potential antioxidant activity with IC₅₀ values of the extract and standard reference being 26.7 µg/ml and 29.3 µg/ml respectively.

The study also reveals that *H. abyssinica* methanolic extract exhibited concentration dependent scavenging activity by inhibiting the DPPH radicals generated. Similar studies on the Sudanese

H. abyssinica ethanolic extract indicate that it has antioxidant activity using superoxide radicals [6].

Phytochemicals/Test	Part of the plant	
	Rhizome	Flowers
Alkaloids		
1. Mayer,s	+	+
2. Drangedorff	+	+
Cardioglycosides (Keller-killan)	+	+
Anthraquinone glycosides		
3. Borntegers	+	+
4. Modified Borntegers	+	+
Saponins	-	-
Tannins	+	+
Phenols	+	+
Steroids	+	+
Coumarins	-	-
Flavonoids	+	+
Volatile oils	-	-
Fatty acids	+	+
Terpenoids	+	+


Table 1: Phytochemical tests of various parts of *Hydnora abyssinica*

4. Conclusion and Recommendation

In the light of the above results, phytochemical analysis results are important in the authentication of plant and further processing for isolation of active compounds from the plant. It can be predicted that the antioxidant activity may be due to the presence of phenolics. The findings that this plant has many a wide range of phytochemicals indicate that it may be explored for principles that may supply lead molecules. Keeping in view of

this work, then suggestions for carrying out further studies on bioassay guided isolation especially using the and methanolic extracts to obtain and characterize potential antioxidant compounds may be warranted.

Graph 1: In vitro antioxidant Assay (DPPH Assay)



IC₅₀ HA-MeOH Extract-26.7

IC₅₀ ASA-Std 29.3

Acknowledgement

The author greatly acknowledges Mount Kenya University for the Vice Chancellor's Research and Innovation grant that enabled this work this far. Much more credit also is for Dr. Peter Kirira of Mount Kenya University Research and Innovation center for the provision of chemical reagents and review of this work.

References

- Ahmad Sayeed (2007). Pharmacognosy, Introduction of plant constituents and their tests. Jamia Hamdard. New Delhi-110062
- Chang H., Ho Y., Sheu M., Lin Y., Tseng S., Huang G. and Chang Y. (2007). Antioxidant and free radical scavenging activities of *Phellinus merrilli* extracts. *Botanical studies*. 48:407-417.
- Evans W.C. Trease and Evans (2008). Pharmacognosy 15th Ed. WB Saunders Company Ltd., London. p. 241-243.
- Ganesan S. and Roopam Y. Bhatt. (2008). Qualitative nature of some traditional crude drugs available in commercial markets of Mumbai, Maharashtra, India. *Ethnobotanical leaflets*. 12: 348-360.
- Houghton Peter J. and Raman Amala. Laboratory Handbook for the fractionation of Natural Extracts Pharmacognosy Research Laboratories. Chapman and Hall. London. 1998, p. 154-188.
- Koko S.W., Osman E.E. and Galal M. (2009). Antioxidant and antiglycation potential of some Sudanese medicinal plants and their isolated compounds, *Bioletín Latinoamericano y del Caribe de Plantas Medicinales y Aromáticas*, 8(5):402-411
- Kokwaro J.O. (2009). Medicinal plants of East Africa (Third edition) Kenya: East Africa Literature Bureau.
- Maundu M.K. (1999). Traditional food plants of Kenya. Kenya: National Museums of Kenya.
- Musila W., Kisangau D. and Muema J. (2004). Conservation status and use of medicinal plants by traditional medical practitioners in Machakos District, Kenya. National Museums of Kenya. Available at <http://www.ed.psu.edu/ick2004proceedings/Accessed 18-11-2013>.
- Musselman J.L. and Visser J.H (1987) *Hydnora johannis* in Southern Africa. *Dinteria NR. 19 - Windhoek S.E.A. -MAI*. 77-82.
- Ndwigah S. N. (2013). Phytochemical, antibacterial and antifungal study of *Dombeyatorrid* (J.F. Gmel) and *Hydnora abyssinica* (A. Braun). PhD Thesis. University of Nairobi.
- Nickaval B., Kamalinejad M. and Izadpanah H. (2007). Invitro free radical scavenging activity of five *Salvia* species. *Pak. J. Pharm. Sci.*, 20 (4), 291-294
- Ogochukwu C.S., Uche A. and Ifeanyi O. (2013). Preliminary phytoscreening of different solvent extracts from stem bark and roots of *Dennetia tripetala* G. Baker. *Asian Journal of Plant science and Research*, 3(3):10-13.
- Ruiz-Terán F., Medrano-Martínez A. and Navarro-Ocaña (2008). Antioxidant and free radical scavenging activities of plant used extracts used in traditional

- medicine in Mexico. *Afr. J. Biotechnol.* 7(12):1886-1893.
15. Saadabi A.M.A and Ayoub S.M.H. (2009). Comparative bioactivity of *Hydnora abyssinica* A. Braun against different groups of fungi and bacteria. *Journal of Medicinal Plants Research.* 3(4) 262-265.
 16. Savithrama N., Linga Rao M. and Suhrulatha D. (2011). Screening of Medicinal Plants for secondary metabolites. *Middle East J. Sci. Res.*, 8(3):579-584.
 17. Srivastava P., Raut N. H., Puntambekar M. H., Upadhye S. A. and Desai C. A (2010). Stability studies of crude plant material of *Bacopa monnieri* and its effect on free radical scavenging activity. *J Phytol* (2010) 103-109.
 18. Tennakoon K.U., Bolin J .F. Musselman J.A. and Maass E. (2007). Structural attributes of the Hypogeous holoparasite *Hydnora triceps* Drège and Meyer (Hydnoraceae). *AJB.*
 19. Yadav RNS. and Agarwala (2011). Phytochemical analysis of some medicinal plants. *Journal of Phytology*, 3(12): 10-14.