

Studies on the effects of stinging nettle extract, phosphoric acid and conventional fungicide combinations on the management of potato late blight and tuber yield in the Highlands of Kenya

Direct Research Journal of Agriculture and Food Science (DRJAFS) Vol.2 (8), pp. 119-127, September 2014

Available online at directresearchpublisher.org/drjafs

ISSN 2354-4147 ©2014 Direct Research Journals Publisher

Research Paper

M. Njogu¹, R. Nyankanga², J. Muthomi² and E. Muindi³

¹Chuka University, P.O BOX 109-60400 Chuka, Kenya.

²University of Nairobi, P.o Box 29053, Kangemi-Kenya.

³Sokoine University of Agriculture, P. o. Box 3008, Morogoro-Tanzania.

*Corresponding Author E-mail: mrtnnjogu@gmail.com; Tel: +2540734617131; Fax: 0202310302

Accepted 12 August, 2014

ABSTRACT

Efficacy of stinging nettle extract, phosphoric acid and fungicides combinations was assessed for late blight control for two cropping seasons at Tigoni and Marimba locations in Kenya. Disease severity, relative area under disease progress curves (RAUDPC), pathogen lesions sizes and numbers, and tuber yield were quantified during the 2008 and 2009 cropping cycles. The application of metalaxyl alternated with phosphoric acid resulted in significant suppressive effects on late blight. The average late blight severity ranged from 3.5 to 34% in 2008 and 4.7 to 50% in 2009 at Tigoni location. RAUDPC for the same location ranged from 5 to 40% and 5 to 50% in 2008 and 2009, respectively. Similar levels of late blight severity have been recorded at Marimba locations. Lesion, growth/size and pathogen lesion numbers on potato plants differed significantly ($p < 0.05$) among treatments. However,

stinging nettle extract, phosphoric acid and fungicides varied in late blight control. Potato tuber yield varied among treatments. Phosphoric acid treatment had significantly ($p < 0.05$) greater tuber yield compared to those of metalaxyl at both locations. Field plots treated with stinging nettle extracts resulted in the lowest tuber yield than the corresponding treatments with the exception of the untreated control. Thus, stinging nettle extract, phosphoric acid, fungicides and their combinations can be readily effective in the suppression of late blight severity and pathogen lesions with moderate increases in tuber yield.

Key words: Stinging nettle extract; phosphoric acid; fungicide combinations; late blight; potato yield.

INTRODUCTION

Management of potato late blight caused by *Phytophthora spp.* remains a major challenge worldwide, especially among small-scale farmers in the tropical highlands. In Kenya, attempts to control late blight are almost entirely accomplished by the use of conventional fungicides like Borduo mixture and cultivars with low to moderate levels of resistance (Haverkort 1990). In most cases, there is an inadequate control of the disease resulting in high costs and complete crop failure.

It has been estimated that 30–60% of the crop is lost due

to late blight annually in Kenya (Nyankanga *et al.*, 2004). Metalaxyl and mefenoxam have been effective in controlling late blight previously, but metalaxyl-resistant populations of *P. infestans* have been reported in many parts of the world (Dowley and O'Sullivan 1981; Derie and Inglis 2001).

The use of alternative products or combinations of products can reduce fungicide inputs and enhance integrated crop management (Cao and van Bruggen 2001). Some chemical compounds have been shown to increase natural plant defense mechanisms (Gozzo 2003). These chemical

compounds can be used as components of integrated pest management. Amino butyric acid and fosetyl-Al applied to foliage at early stages of crop growth can increase the resistance of potato foliage and tubers to late blight (Andreu *et al.* 2006). Phosphoric acid has been reported to inhibit sporulation of the pathogen and adversely impact pathogen infection (Schwinn and Margot 1991). Both phosphoric acids and fosetyl-Al have been reported as effective in controlling plant diseases caused by oomycetes (Coffey and Bower 1984; Ouimette and Coffey 1989b). The mode of action of phosphonate has been reported to be direct anti-fungal activity by preventing mycelia growth (Fenn and Coffey 1989) and indirect stimulation of host defenses (Guest and Grant 1991). These compounds are also thought to have some systemic characteristics with basipetal and acropetal activity for control of foliar and tuber blight (Cooke and Little 2001). This chemical characteristic allows the compounds to be stable in plants for several weeks and offer long-term protection in potato leaves and tubers (Ouimette and Coffey 1989a; Smillie *et al.*, 1989).

To the best of our knowledge, there is no published research detailing the effect of stinging nettle plant extract on potato late blight control. However, plant extract from stinging nettle has been reported to be effective against a different pathosystem, *Phomopsis theae* of tea (Onyango *et al.*, 2005). Plant extract from stinging nettle, phosphoric acid and fungicide combinations could offer a low-cost strategy for managing potato late blight in a region where disease occurrences are prevalent. Therefore, the objective of this study was to determine the efficacy of phosphoric acid, plant extract from stinging nettle, fungicides and their combinations for the management of potato late blight in comparison with fungicides such as premix of Ridomil® (metalaxyl and mancozeb) and Mancozeb® currently used for late blight management in this region.

MATERIALS AND METHODS

Experimental design and layout

The experiment was carried out at the National Potato Research Centre (NPRC), Tigoni and Marimba location in Meru. The two sites have different climatic conditions of (temperature, relative humidity and rainfall) and late blight pressure (Ojiambo *et al.*, 2001). Late blight is a perennial problem at these locations. Treatments were laid in a split plot, in a randomized complete block with three replicates. The main plot consisted of stinging nettle extract and fungicide spray treatments, while the subplots had two potato cultivars grown in many potato cultivation areas of Kenya. Desiree, a cultivar susceptible to late blight and grown in many potato cultivation areas of Kenya for fresh tuber consumption, and Tigoni, a late blight-resistant cultivar grown in many potato production regions for making chips, were used. The main plots were 4.5 x 3.0 m (length x

width) while the subplots were 3.0 x 2.25 m (length x width). Potatoes were planted in furrow at a spacing of 0.75 m between rows and 0.3 m within rows. The distance between blocks was 2 m wide, while an alley of 1.5 m separated the subplots. Two outer rows were used as guard rows while disease severity was recorded in two middle rows and a border of 1 m width surrounded each potato plot. Standard agronomic practices for this region were used for all trials. Sprouted certified potato seeds were planted on 23 October and 11 November 2008 at Tigoni and Marimba, respectively, during the short rainy season, and 13 March and 28 March 2009 at Tigoni and Marimba, respectively, during long rainy season. Diammonium phosphate (DAP) fertilizer was applied at the rate of 500 kg/ha. Field plots were hilled two times during crop growth at both locations. Supplemental irrigation was applied during dry periods as needed. Late blight development was initiated by natural inoculum present in the field sites during the cropping cycle.

Treatment applications

Seven treatments were applied for control of late blight as follows: (1) a premix of Ridomil (64% mancozeb and 4% metalaxyl) at the rate of 2.5 kg/ha, (2) Dithane M45 (80% mancozeb) at the rate of 2.5 kg ha⁻¹, (3) (mono dipotassium) at the rate of 2.5 L ha⁻¹, (4) extract of stinging nettle at the rate of 50 g/ L, (5) mancozeb alternated with phosphoric acid, (6) Ridomil followed by application of phosphoric acid in subsequent application and (7) control treatment. Mancozeb and the combination of mancozeb and metalaxyl (Ridomil) were used as the protective and curative fungicide treatments, respectively. Fungicides were applied at 7-day intervals except for mancozeb and metalaxyl which were applied two times during the cropping season. The sprayer was calibrated prior to commencement of applications so as to deliver spray volume of 400 L/ha at a pressure of 3 bars. The potato plants in the untreated plots were sprayed with water. Treatment applications were initiated at the onset of late blight symptoms initiated from natural inoculum. To prepare plant extract, stinging nettle plants were harvested from the University of Nairobi Farm located at Kanyariri during the flowering stage. Plant extract was prepared by crushing naturally dried whole plants except the roots into fine powder. The powder extract was filtered through a 5-mm mesh sieve to obtain a fine powder. Fifty grams of the fine powder was soaked in 1 litre of water for 24 hours and then added to 20 litres of water for application to foliage.

Late blight assessment

Late blight parameters assessed consisted of disease severity, disease incidence, lesion size, number of lesions

Table 1. Effects of stinging nettle extract, phosphoric acid and Fungicide combinations on average disease severity (%) and relative area under disease progress curves (RAUDPC) on two potato cultivars at Tigoni location during 2008 and 2009 seasons.

Treatments	Tigoni location 2008				Tigoni location 2009			
	Disease severity		RAUDPC		Disease severity		RAUDPC	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	15.1	8	18.4	9.8	28.4	12.7	28.2	13.1
Phosphoric acid (Phosphite)	9	4.4	11.3	5.6	10	5.9	10.3	5.8
Mancozeb + metalaxyl(Ridomil)	10.2	3.9	13	4.9	10.9	5.5	11.2	5.6
Macozeb (Dithane M45)	12.8	5.87	15.5	7.5	27.6	12.5	27.1	12.6
Ridomil + Phosphoric acid	6.9	3.5	8.9	4.6	8.8	4.7	9.1	4.8
Mancozeb + phosphoric acid	10.2	4.4	12.8	4.6	13.8	7.1	14.1	7.1
Untreated	33.6	13.6	39.4	16.5	50	22.8	50.2	22.8
LSD ($P = .05$) Treatment	2.7	1.7	6.6	3.2	2.4	2.1	2.6	2.2
LSD ($P = .05$) Variety	0.7	0.4	2.7	2.2	1.3	1.1	1.6	1.3
% CV	9		7.1		2		3.1	

and tuber blight incidence. Field plots were assessed for late blight severity by visual rating based on percent leaf area blighted using a scale of 0– 100%, where 0% refer to no disease and 100% refers to total foliage damage (James 1971). Assessment for late blight was done in the two inner rows of a 4-row plot beginning from the time when symptoms were first observed. Subsequent disease assessments were made at 4-day intervals until the potato foliage in the untreated plots was completely destroyed. The size of lesions was determined from five lesions on each plant for a total sample size of five plants per subplot. Lesion measurements were repeated on the same plants on 4-day intervals until the end of the season. The number of lesions per plant was determined in five plants per subplot.

Tuber blight assessment

At physiological maturity, tubers from each of the 10 hills per sub-plot were harvested separately and diseased tubers were identified visually and tuber numbers were counted. Tubers with external symptomatic lesions typical of blight were sliced to confirm the presence of brown discoloration backed up by pathogen isolation. Tuber blight incidence was calculated by counting the number of diseased tubers and expressing it as a percentage of the total number of tubers per plot.

Effect of fungicide on potato tuber yield and yield components

At physiological maturity, plants were dehaulmed 2 weeks prior to harvesting. At harvest, tubers from 10 hills in the middle of the two inner rows of each plot were harvested by carefully lifting out the tubers. The tubers were graded

according to size as follows: chatt (<25 mm), seed (25–30 mm) and ware (>55 mm) in diameter. Tubers were also separated into marketable (>6.4 cm diameter) and unmarketable (<6.4 cm diameter) and subsequently weighed separately. The number of tubers per hill was also recorded and yield was converted to tonnes per hectare.

Statistical analyses

The area under disease progress curve (AUDPC) was computed from disease severity using the method previously described (Shaner and Finney 1977). Disease and yield data with the exception of lesion expansion rate were subjected to analysis of variance (ANOVA) using the PROC ANOVA procedure of Genstat (LAWES Agricultural Trust Rothamsted Experimental station 2006, version 9). The lesion expansion rate data were analysed using repeated measures analysis, and differences among the treatment means were compared using the Fisher's protected least significant difference (LSD) test at 5% significant level.

RESULTS

Effect of fungicides, phosphate and plant extract on late blight severity

Plots treated with mancozeb had a disease severity in the range of 12.8–27.6% compared to those plants treated with extract from stinging nettle, which had disease severity of 15.1–28.4% on Desiree, the susceptible cultivar (Table 1). However, there were no significant differences between the two treatments on the resistant cultivar. There were no significant differences between mancozeb phosphoric acid and

Table 2. Effects of fungicide and stinging nettle extract combinations on mean disease severity (%) and relative area under disease progress curve (RAUDPC) on two potato cultivars at Marimba location in 2008 and 2009 seasons.

Treatments	Marimba location 2008				Marimba location 2009			
	Disease severity		RAUDPC		Disease severity		RAUDPC	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	17.8	9.1	21.5	14.3	30.5	13.7	30.4	13.9
Phosphoric acid	10.2	4.6	16.2	5.8	11	6.2	11.4	6.4
Mancozeb + metalaxyl	10.4	4.4	16.4	5.5	12.1	6.2	12.4	6.3
Mancozeb (Dithane M45)	14	6.3	17.1	11.1	29.1	13.3	24.7	12.9
Ridomil + Phosphoric acid	7.4	3.7	12.7	4.7	9.6	5.3	10	5.5
Mancozeb + phosphate	10.8	5.4	16.4	6.9	15.8	8.1	16.4	8.1
Untreated	40.1	18	49.1	24.3	53.2	24.5	52.5	24.1
LSD ($P = .05$) Treatment	2.5	2.3	8.4	5.2	2.8	2.2	3.1	2.7
LSD ($P = .05$) Variety	0.94	0.8	5.1	4.7	1.4	1.1	1.1	1.2
% CV	8.5		8.1		0.4		1.3	

Table 3. Mean *Phytophthora infestans* lesion growth rate (mm/day) on two potato cultivars at Tigoni and Marimba locations during 2008 and 2009 cropping seasons.

Treatments	Tigoni 2008		Tigoni 2009		Marimba 2008		Marimba 2009	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	4	2.9	4.3	3.2	4.3	3	4.4	3.1
Phosphoric acid	2.7	2.1	2.8	2.5	2.8	2.4	2.8	2.4
Mancozeb + metalaxyl	2.5	2.3	2.8	2.4	2.8	2.3	2.9	2.4
Mancozeb (Dithane M45)	3.4	2.3	5.7	2.9	3.7	2.7	4.3	3
Ridomil + Phosphoric acid	1.9	1.4	2.4	2.1	2.5	1.2	2.2	2.1
Mancozeb + phosphoric acid	2.7	2.2	4.2	2.2	2.8	2.5	3.4	2.7
Untreated	6.8	3.8	8	4.4	9.2	6.2	8.4	4.5
LSD ($P = .05$) Treatment	0.9	0.7	3	0.5	0.6	0.5	0.5	0.4
LSD ($P = .05$) Variety	0.5	0.4	1.5	0.3	0.33	0.4	0.4	0.3
% CV	2.9		2.9		1.9		0.8	

mancozeb-treated plots for both potato cultivars at Tigoni site in 2008, but in 2009, the untreated plots had significantly ($P = .05$) higher late blight severity and RAUDC than the rest of the treatments (Table 1). The application of metalaxyl and mancozeb followed by phosphite treatment had the least level of late blight disease at both locations (Tables 1 and 2). In 2008 cropping season, there was considerably lower disease levels at Tigoni than at Marimba. Late blight severity was significantly ($P = .05$) higher in 2009 compared to 2008 at the Tigoni location.

Effect of stinging nettle extract, phosphoric acid and conventional fungicides on lesion size and numbers

Mean lesion expansion increased on both cultivars and at

both locations in all the plots during the 2 years of the experiment (Tables 3 and 4). There was a variation in lesion expansion rates in plots treated with fungicide combinations and plant extract. Potato foliage treated with metalaxyl + mancozeb (2.5 kg/ ha) followed by phosphoric acid (2.5 L/ ha) had lower lesion sizes than the other treatments. At both sites, the use of phosphoric acid in combination with either metalaxyl + mancozeb (Ridomil) or mancozeb did not lead to any significant reductions in lesion size expansion on the cultivar Tigoni (Table 3). The extract from stinging nettle reduced pathogen lesion size significantly ($P = .05$) compared to the untreated plots. The plant extract from stinging nettle and mancozeb treatment resulted in similar lesion expansion rates at Tigoni location (Table 3). Lesion expansion rate on the cultivar Desiree was significantly greater than on Tigoni cultivar at both sites.

Pathogen lesion numbers quantified at both sites

Table 4. Effects of stinging nettle extract, phosphoric acid and fungicides combinations on average number of lesions per plant quantified on two potato cultivars at Tigoni and Marimba locations in 2008 and 2009 cropping seasons.

Treatments	Tigoni 2008		Tigoni 2009		Marimba 2008		Marimba 2009	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	11.1	8	16.7	8.8	13.4	8	17.4	9.1
Phosphoric acid	5.6	4	7.2	4.6	6.8	3.3	7.8	4.9
Mancozeb + metalaxyl	6.1	4	7.5	4.7	6.5	3.7	8	4.8
Mancozeb (Dithane M45)	9.2	5	15.5	8.3	10.4	5.3	16.1	8.6
Ridomil + Phosphoric acid	4.4	3	5.9	3.9	5	3.4	6.3	4.1
Mancozeb + phosphoric acid	8.3	4	10.6	6.3	8.9	4.8	11.3	6.8
Untreated	21.8	12	29.1	16.2	23.8	12.1	30	16.9
LSD ($P = .05$) Treatment	2.3	2.1	2.1	1.7	1.8	1.4	1.8	1.6
LSD ($P = .05$) Variety	0.6	0.4	0.4	0.3	0.6	0.7	0.4	0.4
% CV	8.5		10.1		7		6.9	

Table 5. Incidence of tuber blight (%) on two potato cultivars treated with stinging nettle extract, phosphoric acid, different fungicides, and fungicides combinations during the 2008 and 2009 cropping seasons at Tigoni and Marimba locations.

Treatments	Tigoni 2008		Marimba 2008		Tigoni 2009		Marimba 2009	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	4	7.3	2.1	4.8	1.2	2.4	1.3	2.9
Phosphoric acid	1.4	4.3	1.1	1.1	0.7	1.4	0.7	1.1
Mancozeb + metalaxyl	1.4	5.2	1.4	1.3	0.8	2.6	0.7	1.5
Mancozeb (Dithane M45)	3.2	7.4	1.8	3.1	1.4	3.8	1.7	2.9
Ridomil + Phosphoric acid	0.5	3.3	0.8	0.8	0.5	1.3	0.3	1
Mancozeb + phosphoric acid	0.7	2.8	1.5	2.5	0.9	2.6	1.1	1.9
Untreated	0.9	7.4	1.9	3.4	1.3	3.5	0.8	1.7
LSD ($P = .05$) Treatment	2.5	2.1	2.1	0.3	1.4	1.2	0.9	0.7
LSD ($P = .05$) Variety	1	0.7	0.5	0.3	0.2	0.32	0.3	0.16
CV (%)	26		36.4		11.6		8.7	

were generally low at 55–61 days after planting for all treatments but increased rapidly in plots treated with extract from stinging nettle and in the control (unsprayed) treatment. Mean lesion numbers in all treatments differed significantly ($P = .05$) with the control treatments on both cultivars at the two sites (Tables 3 and 4). Treatments of metalaxyl + mancozeb (Ridomil) followed by phosphite application resulted in the least number of pathogen lesions. Variation in lesion numbers was recorded in various treatment combinations of mancozeb (2.5 kg ha⁻¹), phosphite, and extract of stinging nettle extract on the cultivar Desiree. However, on the cultivar Tigoni, treatment sequence of metalaxyl + mancozeb followed by phosphite did not differ with other fungicides treatments with the exception of plant extract from stinging nettle (Table 4). At Marimba location, a similar trend was observed where

treatments of metalaxyl + mancozeb followed by phosphite resulted in the lowest number of pathogen lesions on both cultivars. Lesion numbers were similar in field plots treated with either metalaxyl + mancozeb or phosphite by itself on both cultivars (Table 4). The lesion numbers were reduced in treatment of extract of stinging nettle compared to the untreated plot.

Effects of stinging nettle extract, phosphoric acid and fungicides on tuber blight incidence

Tuber blight incidence was observed at harvest in all the treatments and at both locations (Table 5). The blight incidence on tubers was in the range of 0.7–7.4% on both cultivars during 2008. There were significant ($p < 0.05$) differences between the two

Table 6. Effects of stinging nettle extract, phosphoric acid, fungicides and treatment combinations on total yield of two potato cultivars during 2008 and 2009 cropping years.

Treatments	Total yield (t/ha)– 2008				Total yield (t/ha)– 2009			
	Tigoni location		Marimba location		Tigoni location		Marimba location	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	7.7	12	7.6	8.9	11	15.3	8.1	9.7
Phosphoric acid	15.6	17	14.2	18	15.6	24	13.7	16.2
Mancozeb + metalaxyl	19.2	16	20.1	16.4	14.4	22.5	17.1	19.6
Macozeb (Dithane M45)	9.1	13.5	8.8	9.7	9	15.3	8.6	9.9
Ridomil + Phosphoric acid	18.3	21	20.1	17.8	18.6	26.47	22.7	21.9
Mancozeb + phosphoric acid	11.8	16	8.8	11.1	15.7	21.5	8.9	10.7
Untreated	4.8	6.6	5.3	5.8	7.5	9.9	5.6	6
LSD ($P = .05$) Treatment	6.8	5.8	5.9	4.8	6.7	5.2	4.5	3.6
LSD ($P = .05$) Variety	0.7	0.6	1.17	1.3	2.5	1.9	1.1	0.8
CV (%)	17.1		26.3		16.5		10.7	

Table 7. Effects of stinging nettle extract, phosphoric acid fungicides, and treatment combinations on marketable yield on two potato cultivars during 2008 and 2009 cropping years.

Treatments	Marketable yield t/ha in 2008				Marketable yield t/ha in 2009			
	Tigoni location		Marimba location		Tigoni location		Marimba location	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	9.4	10.8	7.5	8.9	10.1	13.3	6.9	8.6
Phosphoric acid	13.8	16.8	13.4	17	14.4	23.3	13.4	15.3
Mancozeb + metalaxyl	16.9	13.4	18.4	15.5	13.5	20.8	16.1	18.3
Macozeb (Dithane M45)	5.1	11.8	6.6	7.9	7.9	14	7.5	8.9
Ridomil + Phosphoric acid	14.2	18.5	18.6	16.8	17.5	25.7	21.4	20.9
Mancozeb + phosphoric acid	10.2	14.4	8.4	10	14.7	20.4	8.5	9.7
Untreated	3.9	4.8	3.8	4.3	6.9	8	3.9	4.4
LSD ($P = .05$) Treatment	5.2	4.8	6.8	4.9	6.8	4.9	4.4	3.7
LSD ($P = .05$) Variety	1.6	1.6	2.6	2.2	2.6	2.2	0.9	0.6
CV (%)	26.6		18		18.5		10	

potato cultivars in tuber blight incidence at harvest (Table 5). In 2008, tuber blight incidence on cultivar Desiree in the untreated plots differed significantly ($P = .05$) with the rest of the treatments (Table 5). On the cultivar Tigoni, tuber blight incidence was similar in plots treated with mancozeb, extract from stinging nettle and the untreated control. In 2009, the incidence of tuber blight ranged from 0.4 to 3.8% across cultivars at Tigoni location.

Effect of stinging nettle extract, phosphoric acid and fungicides on tuber yield Parameters

Tuber yield varied between cultivars and years. At the Tigoni location, total tuber yield ranged from 4.8 to 19.2 and 6.6 to 21 t/ha on Desiree and Tigoni cultivars, respectively,

during 2008 (Table 6). At Marimba location, the range in tuber yield was 5.3–20.1 on Desiree and 5.8–17.8 t/ha on the cultivar Tigoni (Table 6). In 2009, tuber yield at Tigoni ranged from 7.5 to 18.6 t/ha on Desiree and 9.9–26.7 t/ha on cultivar Tigoni (Table 6). Tuber yield was generally greater at Tigoni than the Marimba location. At both locations, the highest tuber yield was recorded on treatments where metalaxyl + mancozeb was followed by phosphite application. The untreated control plots had the lowest tuber yield recorded during both years. Stinging nettle extract did not result in any significant increases in tuber yield compared to the untreated control on the cultivar Tigoni but resulted in significantly higher yield on cultivar Desiree (Table 6). All the treatments resulted in significantly ($P = .05$) greater marketable yield compared to the untreated plots where no fungicides or other treatments were applied (Table 7). The marketable yield generally ranged from 3.9

Table 8. Effects of stinging nettle plant extract, phosphoric acid, fungicides, and treatment combinations on tuber numbers on two potato cultivars during 2008 and 2009 cropping years.

Treatments	Tuber numbers/hill 2008				Tuber numbers/hill 2009			
	Tigoni location		Marimba location		Tigoni location		Marimba location	
	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety	Desiree variety	Tigoni variety
Stinging nettle extract	5.8	8	6.4	6.2	6.3	8	5.9	6.5
Phosphoric acid	7.7	10	8.33	8.7	9.7	14	8.3	10.7
Mancozeb + metalaxyl	7.5	11	6.3	9.3a	9.3	13	7.7	11.3
Mancozeb (Dithane M45)	4.9	8	4.7	6.8	6.7	8.3	6	8.3
Ridomil + Phosphoric acid	7.1	12	8.4	12.3	10.7	14.3	10.3	13.7
Mancozeb + phosphoric acid	6.9	8	5.9	6.6	7.7	10.7	6.3	7.7
Untreated	4.6	6	3.3	4	5.3	6.3	4	4.3
LSD ($P = .05$) Treatment	1.6	1.4	1.7	1.6	1.4	1.2	4.5	3.6
LSD ($P = .05$) Variety	0.6	0.4	0.8	0.6	0.7	0.7	1.1	0.8
CV (%)	6.62		5.8		10.6		10.7	

PI. reduce table numbers.

to 16.8 t/ha across years and cultivars (Table 7). Potato crops in which metalaxyl + mancozeb and metalaxyl + mancozeb followed by phosphite application resulted in the highest marketable tuber yield of 16.93 and 18.53 in Desiree and Tigoni cultivars; respectively (Table 7). Alternating phosphite with mancozeb resulted in significantly higher marketable yield compared to plots where mancozeb alone was applied on Desiree but not on Tigoni cultivar. Similarly, all treatments of mancozeb followed by phosphite; metalaxyl + mancozeb and metalaxyl + mancozeb followed by phosphite and phosphite alone resulted in significantly ($P = .05$) higher number of tubers per hill than the untreated control plants at Tigoni location (Table 8). Similar treatment effects produced significant marketable yield at Marimba location.

DISCUSSION

This study revealed that there was very little variance in the incidence of tuber blight recorded among treatments of fungicides, phosphite, and plant extract combinations at both locations and seasons. The fungicide treatments reduced late blight severity and AUDPC as well as lesion size and numbers relative to the untreated control. Treatments of phosphite also resulted in the reduction of late blight disease on potato as compared to those of untreated control. The extract from stinging nettle, however, had the least suppressive effects on late blight. Efficacy of fungicide control of late blight on potato has been previously documented in tropical environments due to their fungicidal properties (Olanya *et al.*, 2001, 2006; Namanda *et al.*, 2004; Nyankanga *et al.*, 2004). Treatment combinations involving metalaxyl + mancozeb (Ridomil) and phosphite had much more effective late blight control as was evident by the lower late blight damage on treated plants compared to mancozeb or extract of stinging nettle. In general, better absorption of metalaxyl + mancozeb under various

environmental conditions has been shown to be effective for control of most plant diseases incited by oomycetes (Easton and Nagle 2007).

The systemic nature of metalaxyl and its efficacy has been previously documented, particularly in populations of late blight with A1 mating type (Mukalazi *et al.* 2001). The relatively moderate late blight disease recorded in treatment of mancozeb suggests that protectant fungicides may have variable protectant effects on late blight under diverse conditions. In contrast to the above findings, research reports in other highland tropics showed that mancozeb can be effective in reducing the impacts of late blight (Namanda *et al.*, 2004). The differences in disease levels between Tigoni and Marimba may be attributed to difference in weather conditions at these locations. Variation in late blight incidence and severity has been previously reported among the various locations of Kenya (Nyankanga *et al.*, 2004; Olanya *et al.*, 2006). The treatment of metalaxyl + mancozeb and the application of phosphoric acid had the highest suppressive effects on late blight epidemics. The enhanced disease control obtained suggests that this treatment combination (metalaxyl + mancozeb) can be most effective on late blight management. The incorporation of phosphite for late blight management subsequent to mancozeb fungicide also improved the suppressive effect of metalaxyl + mancozeb and mancozeb more than the use of mancozeb by itself on the susceptible potato cultivars.

The fungicides and phosphoric acid treatments had significant effects on pathogen lesion numbers and size compared to those of plant extract from stinging nettle or the untreated control. Although phosphoric acid was reported to be effective in controlling lesion development and sporulation on *P. infestans* in previous studies (Johnson *et al.* 2004), its effects did not differ with that of metalaxyl + mancozeb under the tropical conditions of our experiment conducted in Kenya. Lesion developments also differed between the locations of Tigoni and Marimba,

perhaps due to the cooler environment at Marimba.

In previous research, it was observed that phosphoric acid and other protectant fungicides applied in the field prior to harvest and on harvested tubers can protect potato tubers from *P. infestans* upon subsequent inoculation (Cooke and Little 2001; Johnson et al. 2004). The same studies also demonstrated that foliar applications of phosphoric acid can reduce late blight infections occurring during the growing season (Cooke and Little 2001). Therefore, our findings are in agreement with the results of previous studies. The application of plant extract from stinging nettle did not have any significant suppressive effects on tuber blight incidence on both cultivars at harvest in our experiments and is not totally unexpected. This is similar to other previous findings in which plant extracts such as aerated compost tea and effective micro-organism mix were shown to have very little reduction in late blight disease on potato foliage (Cao and Van Bruggen 2001; Olanya and Larkin 2006; Al-Mughrabi 2007).

Variation in tuber blight incidences and disease control by fungicides in different treatments and locations has been reported in previous studies (Dorrance and Inglis 1998; Platt and Tai 1998; Nyankanga et al., 2007). It has been shown that potato tubers from field plots that are not protected by fungicides often have a lower incidence of tuber blight than do tubers from plots that have been protected due to the presence of suppressive soil conditions and other environmental factors that affect tuber blight development on potato (Andrivon 1995). This finding has been attributed to the fact that late blight epidemics on unprotected plants proceed more rapidly than disease development on fungicide-protected plants, leading to a limited sporangia from lesions which can be washed from the foliage onto tubers (shorter duration of epidemic) than for the unprotected plants. Similarly, the lack of differences in tuber blight incidence between Tigoni (resistant check) and Desiree (susceptible check) cultivars imply that foliar resistance may have limited effect on tuber blight incidence.

Potato tuber yield was significantly impacted by fungicide treatments and phosphite but to a lesser degree with the application of plant extract from stinging nettle in these experiments. Fungicides have shown greater efficacy compared to plant extracts due to their active ingredients which are often more potent for disease control compared to plant extracts. For example, successful fungicide controls for late blight and tuber yield have been documented in various studies (Fontem and Aighewi 1993; Ojiambo et al., 2001; Namanda et al., 2004). This has been attributed to fungicide protection of potato foliage, thereby increasing yield potential in treated plots compared to the untreated control plots. Alternating phosphite with mancozeb for late blight control on foliage resulted in significantly higher tuber yield than the application of mancozeb alone. This indicates that the treatment combinations can result in better blight control and tuber yield. On the contrary, the lack of significant difference in tuber

yield between treatments of stinging nettle extract and the untreated control imply that the plant extract treatment had no effect on tuber yield.

It appears that the applications of fungicides, phosphite and plant extract from stinging nettle resulted in effective management of late blight on potato. The combined use of metalaxyl + mancozeb with phosphate applications provided the best overall disease control. The stinging nettle extract, although did significantly reduce disease parameters relative to the untreated control, was overall the least effective of all treatments tested. Late blight severity, AUDPC, lesion development and pathogen lesion numbers varied between cultivars and were significantly greater on Desiree than on Tigoni cultivar at both locations. Low to moderate levels of late blight were recorded at both locations, and fungicides alone, alternating fungicides + phosphite and phosphite applications produced significantly greater late blight control compared to plant extract or the untreated control at both locations. Little variation in tuber blight incidence was detected among treatments and years; however, late blight incidence on tubers was greater on Tigoni than Desiree cultivar. Most treatments resulted in greater tuber and marketable yield compared to the untreated control.

References

- Al-Mughrabi K (2007). Suppression of *Phytophthora infestans* in potatoes by foliar application of food nutrients and compost tea. *Aust. J. Basic Appl. Sci.* 1:785–792.
- Andreu AB, Guevara MG, Wolski EA, Daleo GR, Caldiz DO (2006). Enhancement of natural disease resistance in potatoes by chemicals. *Pest Manag. Sci.* 62:162–170.
- Andrivon D (1995). Biology, ecology, and epidemiology of the potato late blight pathogen *Phytophthora infestans* in soil. *Phytopathology.* 85:1053–1056.
- Cao K, Van Bruggen AHC (2001). Inhibitory efficacy of several plant extracts and plant products on *Phytophthora infestans*. *J. Agric. Univ. Hebei* 24:90–96.
- Coffey MD, Bower LA (1984). In vitro variability among isolates of eight *Phytophthora* species in response to phosphorous acid. *Phytopathology.* 74:738–742.
- Cooke LR, Little G (2001). The effect of foliar application of phosphonate formulations on the susceptibility of potato tubers to late blight. *Pest Manag. Sci.* 58:17–25.
- Derie ML, Inglis DA (2001). Persistence of complex virulence in population of *Phytophthora infestans* in western Washington. *Phytopathology.* 91:606–612.
- Dowley LJ, O'Sullivan E (1981). Metalaxyl-resistant strains of *Phytophthora infestans* (Mont.) de Bary in Ireland. *Potato Res.* 24:417421.
- Dorrance AE, Inglis DA (1998). Assessment of laboratory methods for evaluating potato tubers for resistance to late blight. *Plant Dis.* 82:442–446.
- Easton GD, Nagle ME (2007). Timing and root absorption affecting efficacy of metalaxyl in controlling *Phytophthora infestans* on potato in Northwestern Washington State. *Plant Dis.* 69:499–500.
- Fenn ME, Coffey MD (1989). Quantification of phosphonate and ethyl phosphonate in tobacco and tomato tissues and significance for the mode of action of two phosphonate fungicides. *Phytopathology.* 79:7682.
- Fontem DA, Aighewi B (1993). Effect of fungicides on late blight control and yield loss of potato in the western highlands of Cameroon. *Int J Pest Manag.* 39:152155.

- Gozzo F (2003). Systemic acquired resistance in crop protection: from nature to a chemical approach. *J Agric Food Chem.* 51:4487–4503.
- Guest D, Grant BR (1991). The complex action of phosphonates as antifungal agents. *Biological Rev.* 66:159187.
- Haverkort AJ (1990). Ecology of potato cropping systems in relation to latitude and altitude. *Agric. Syst.* 32:251272.
- James C (1971). A manual of assessment keys for plant diseases. Canada Department of Agriculture Publication No. 1458. St. Paul, MN: APS Press.
- Johnson DA, Inglis DA, Miller JS (2004). Control of potato tuber rots caused by oomycetes with foliar applications of phosphorous acid. *Plant Dis.* 88:11531159.
- Mukalazi J, Adipala E, Sengooba T, Hakiza JJ, Olanya M, Kidanemariam HM (2001). Metalaxyl resistance, mating type and pathogenicity of *Phytophthora infestans* in Uganda. *Crop Prot.* 20:379–388.
- Namanda S, Olanya OM, Adipala E, Hakiza JJ El-Bedewy R, Baghsari AS, Ewell PT (2004). Fungicide application and host-resistance for potato late blight management: benefits assessment from on-farm studies in S.W. Uganda. *Crop Prot.* 23:10751083.
- Nyankanga RO, Wien HC, Olanya OM, Ojiambo PS (2004). Farmers' cultural practices and management of potato late blight in Kenya highlands: implications for development of integrated disease management. *Int. J. Pest Manag.* 50:135–144.
- Nyankanga RO, Wien HC, Olanya OM, Ojiambo PS (2007). Relationship between late blight (*Phytophthora infestans*) of potato on tuber and foliage, as affected by the disease severity on foliage, cultivar resistance and atmospheric and soil variables. *Can. J. Plant Pathol.* 29:372–387.
- Ojiambo PS, Namanda S, Olanya OM, El-Bedewy R, Hakiza JJ, Adipala E, Forbes G (2001). Impact of fungicide application and late blight development on potato growth parameters and yield in the tropical highlands of Kenya and Uganda. *Afr. Crop Sci. J.* 9:225–233.
- Olanya, OM, El-Bedewy R, Ojiambo PS, Ewell PT, Hakiza JJ (2001). Relationships of fungicide application to late blight development and potato growth parameters in the tropical highlands of Uganda and Kenya. In: *Scientist and farmer: partners in research for the 21st century*. International Potato Center (CIP), Program Report 1999–2000. Lima, Peru: CIP. p. 77–86.
- Olanya OM, Larkin RP (2006). Efficacy of essential oils and biopesticides on *Phytophthora infestans* suppression in laboratory and growth chamber studies. *Biocontrol Sci Technol.* 16:901–917.
- Olanya OM, Ojiambo PS, Nyankanga RO (2006). Dynamics of development of late blight (*Phytophthora infestans*) in potato, and comparative resistance of cultivars in the highland tropics. *Can. J. Plant Pathol.* 28:84–94.
- Onyango MO, Khare KB, Achwanya OS, Otieno W (2005). Efficacy of synthetic and non-synthetic fungicides against stem canker disease of tea. *Afri. Crop Sci J.* 13:209–214.
- Ouimette DG, Coffey MD (1989a). Comparative antifungal activity of four phosphonate compounds against isolates of nine *Phytophthora* species. *Phytopathology.* 79:761–767.
- Ouimette DG, Coffey MD (1989b). Phosphonate levels in avocado (*Persea americana*) seedlings and soil following treatment with fosetyl-Al or potassium phosphonate. *Plant Dis.* 73:212–215.
- Platt HW, Tai G (1998). Relationship between resistance to late blight in potato foliage and tubers of cultivars and breeding selections with different resistance levels. *Am. J. Potato Res.* 75:173–178.
- Shaner G, Finney RE (1977). The effect of nitrogen fertilization on the expression of slow-mildewing resistance in knox wheat. *Phytopathology.* 67:1183–1186.
- Smillie R, Grant BR, Guest D (1989). The mode of action of phosphite: evidence for both direct and indirect action modes of action on three *Phytophthora* spp. in plants. *Phytopathology.* 79:921926.
- Schwinn FJ, Margot P (1991). Control with chemicals. In: Ingram DS, Williams PH, editors. *Advances in plant pathology, Phytophthora infestans, the cause of potato late blight*, Vol 7. San Diego, CA: Academic Press Limited. p. 225–265.