



Contents

- I. Diseases in Seed Crop Production

SEMIS - UOM

SEED ENTERPRISE MANAGEMENT INSTITUTE (SEMIs)

Seed Production Short Course

10th – 15th August 2015

Diseases in Seed Crop Production



Prof. James W. Muthomi

Department of Plant Science and Crop Protection

University of Nairobi


Diseases in seed crop production

Disease	Causal agent
Bean anthracnose	<i>Colletotrichum lindemuthianum</i>
Halo blight (bean)	<i>Pseudomonas savastanoi phaseolicola</i>
Common bacterial blight (bean)	<i>Xanthomonas axonopodis phaseoli</i>
Bean common mosaic	<i>Bean common mosaic virus</i>
Head smut (maize)	<i>Sphacelotheca reiliana</i> , <i>Ustilago maydis</i>
Gray leaf spot (Maize)	<i>Cercospora zea-maydis</i>
Maize leaf blight	<i>Drechslera turcicum</i>
Stalk rot / ear rot (maize)	<i>Fusarium graminearum</i> , <i>F. verticillioides</i> , <i>F. proliferatum</i> , <i>F. subglutinans</i> , <i>Stenocarpella maydis</i>
Bacterial blight (cow pea)	<i>Xanthomonas campestris vignicola</i>
Sclerotinia wilt & head rot (sun flower)	<i>Sclerotinia sclerotiorum</i>
Botrytis head rot (sunflower)	<i>Botrytis cinerea</i>

Diseases in seed crop production



Reduced seedling vigour



Seed discolouration,
Shrivelling, rotting &
reduced size



Reduced seedling vigour



How does seed contamination occur?

Seed contamination or infestation

Pathogen itself or parts of it stick or mix with seeds during:

Harvesting

Extraction

Threshing

Selection

Packing

Accompanying contamination

Physical mixing of the seed with pathogen's propagation organs

- Spores
- Sclerotium
- Nematode's galls
- Contaminated plant parts or soil particles containing pathogens

Location of pathogen in seed

- Infection of the embryo
- Under the seed coat
- In the endosperm or cotyledon
- On the surface of seed

How pathogens infect seed

❑ Systemic Infection of the Seed

- Through flowers, fruits or funiculus
- Through the stigma
- Through the wall of the ovary or immature seed covers
- Through wounds & natural openings


Seed contamination or infestation

- Pathogens that stick to the surface of the seed

Accompanying contamination


- Structures of the pathogens
- Mix with infected plant parts
- Soil

Routes of active seed infection



Routes of active seed infection

A. Penetration through ovary wall



E.g.: *Cladosporium variabile* (spinach),
Botrytis spp. (onion)

From Maude (1996)

B. Systemic infection via vascular system



E.g.: Vascular wilt fungi,
endophytes

C. Penetration through floral parts



E.g.: *Ustilago nuda* (grains)
Cucumber mosaic virus

Diseases in seed crop production

Infected seeds



Maize Lethal Necrosis Disease



Loose smut



Head Smut



Maize leaf blight



Gray leaf spot



Maize rust

Maize

Fusarium stalk rot of maize




Charcoal rot



Diplodia stalk and ear rot of maize

Maize



Ear rot of maize

Maize

Fusarium ear rot



Diplodia



Fusarium ear rot



Trichoderma



Aspergillus ear rot



Gibberella ear rot






Anthrachnose



Helminthosporium
leaf blight



Target spot




Head blight

Smut on wheat ears



Wheat kernels with smut symptoms



Wheat scab on ears



Wheat scab symptoms on kernels






Loose smut



Stem rust




Leaf rust



← Powdery mildew




Barley yellow dwarf →




Rice blast

Rice




Bean anthracnose on pods and leaves

Bean



Angular leaf spot on bean



Sclerotinia on bean stems and pods



Aschochyta leaf spot



Web blight



Bean rust




Root rots




Halo blight on bean



Bean virus diseases



Virus diseases



Bacterial blight



Aschochyta




Cercospora



Cowpea



Rust



Anthracnose



Early leaf spot



late leaf spot



Alternaria leaf spot



Rust

Aspergillus crown rot



Ground nut rosette



Virus diseases






Sclerotinia Head Rot of sunflower



MANAGEMENT OF SEED-BORNE DISEASES



Previous cropping

- ❑ Seed production fields should be free from volunteer plants to avoid contamination of the crop seed by:
 - Any seed which is difficult to remove from the crop seed
 - Cross-pollination;
 - Seed-borne diseases transmitted from volunteer plants
 - The previous cropping shall be such that there is the least
 - possible risk of any soil borne diseases being present which
 - could subsequently be transmitted in the harvested seed.

Production in disease-free areas

- Dry areas with low humidity (use irrigation)
- Bean anthracnose and Bacterial blights of bean
- Altering time of planting
- Crop isolation from other fields containing possibly diseased plants

Good production practises

- Use of certified seed
- Minimize plant stress – fertilization & watering
- Weed management
- Well-drained soils
- Seed rate – proper plant density to promote rapid drying of foliage

Eradicate disease-causing pathogen from production area

- Remove alternate hosts and volunteer host plants
- Crop rotation
- Sanitation – residue management
- Creating conditions unfavourable to pathogens
- Polyethylene mulching
- Drip irrigation instead of overhead irrigation
- Soil sterilization for greenhouse & nursery plants
- Seed treatment

Sanitation

- Destroy/ plough under crop residues
- Proper crop handling (wash hands & implements)
- Removal of infected plants (roguing)
- Avoid working in field when wet



Protect crop from disease

- Use resistant/ tolerant crop varieties
- Use of disease-free planting materials
- Spray protective fungicides,
- Protect from vectors
- Control of Insect Vectors

Isolation and Field Inspection

- Seed crops should be isolated from all sources of pollen contamination and seed-borne diseases (including seed-borne virus infection and wild plants that might serve as a source of disease)
- Crop should be inspected at least once at appropriate stage of growth
- At least 20% of the crop of Certified Seed should be inspected
- Presence of any seed-borne disease should be at the lowest possible level

Seed health testing



Germination test



Diseases in seed crop production




Seed health test
for seedborne
pathogens



Fast green test
for physical damage



Tetrazolium test
for seed viability



Tolerated levels for seed borne diseases

Disease	Tolerance level
Head smut (maize)	2 plants per hectare
Loose smut (maize)	2 plants per hectare
Bunt (wheat)	1 head per 100 sq. m
Bunt (sorghum)	1 plant per 1,000 plants
Halo bight (bean)	None at inspection
Anthracnose (bean)	None at inspection
Common bacterial blight (bean)	None at inspection
Bean common mosaic	None at inspection
Bacterial blight (cow pea)	None at inspection
Botrytis head rot (sun flower)	5 plants per 1,000 plants
Sclerotinia wilt & head rot (sun flower)	5 plants per 1,000 plants

THANK YOU